

IOP Conference Series: Earth and Environmental Science



The open access *IOP Conference Series: Earth and Environmental Science (EES)* provides a fast, versatile and cost-effective proceedings publication service.

Latest published conferences

Vol 544

Go

Conference archive

2020

Go

[View forthcoming volumes](#) accepted for publication.

[RSS feed](#)

[Sign up for new issue notifications](#)

JOURNAL LINKS

[Journal home](#)

[Information for organizers](#)

[Information for authors](#)


[Search for published proceedings](#)

[Contact us](#)

[Reprint services from Curran Associates](#)

If you would like more information regarding *IOP Conference Series: Earth and Environmental Science* please visit conferenceseries.iop.org, and if you are interested in publishing a proceedings with IOP Conference Series please visit our page for [conference organizers](#).

- **Conference organizers** can use our [online form](#) and we will get in touch with a quote and further details.
- **Researchers** will enjoy the [conference-based search system](#) to quickly find and browse proceedings of interest. Search through all proceedings by conference title, subject area and conference date / location.



IOP Conference Series: Earth and Environmental Science


Country	United Kingdom -  SIR Ranking of United Kingdom
Subject Area and Category	Earth and Planetary Sciences Earth and Planetary Sciences (miscellaneous) Environmental Science Environmental Science (miscellaneous)
Publisher	IOP Publishing Ltd.
Publication type	Conferences and Proceedings
ISSN	17551315, 17551307
Coverage	2010-2020
Scope	The open access IOP Conference Series: Earth and Environmental Science (EES) provides a fast, versatile and cost-effective proceedings publication service.
	Homepage How to publish in this journal Contact
	Join the conversation about this journal

18


H Index




External Cites per Doc Cites per Doc




Citable documents Non-citable documents



% International Collaboration



Cited documents Uncited documents



Cites / Doc. (4 years)
Cites / Doc. (3 years)
Cites / Doc. (2 years)

IOP Conference Series: Earth and Environmental...

Not yet assigned quartile

SJR 2019
0.18

powered by scimagojr.com

Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scima
```

PAPER • OPEN ACCESS

Metagenomic analysis of diversity and composition of soil bacteria under intercropping system *Hevea brasiliensis* and *Canna indica*

To cite this article: Y Effendi *et al* 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **391** 012023

View the [article online](#) for updates and enhancements.

Metagenomic analysis of diversity and composition of soil bacteria under intercropping system *Hevea brasiliensis* and *Canna indica*

EFFENDI, Y¹, A. PAMBUDI¹, Y. SASAERILA¹, r. S. WijiHastuti¹

¹Department of Biology, Universitas Al Azhar Indonesia. Jl. Sisingamangaraja No. 2, Jakarta Selatan 12110, Jakarta, Indonesia,

Corresponding author: effendiy@uai.ac.id

Abstract. Composition of soil microbiomes plays important ecological roles and vital ecosystem processes in nature of soil. Indeed, interaction between plant root, microbes, and soil influences significantly biologically, physically and chemically of soil properties. In this study, high throughput sequencing of 16S rRNA gene of soil bacteria using Illumina platform was performed for analyzing composition, richness and biodiversity of soil bacteria in two different soil conditions. Two soil samples from Rubber tree – *Canna indica* intercropping areas and two soil samples from area without *C. indica* were collected and analyzed the diversity and richness of soil bacterial communities. Bioinformatic data analysis showed in average more than 5500 bacterial OTU were identified in each soil samples. Relative abundance of individual OTU between soil with and without *C. indica* showed statistically no difference. From 30 phyla that have been identified in both soil conditions, only 5 phyla (BH180.139, Cyanobacteria, Proteobacteria, TM6, and WS3) showed a statistically different ($p < 0.05$) in their abundance. A total 270 and 355 Familia and Genus respectively were identified. There is no statistically difference in richness, abundance and diversity of species between both soil conditions. However, intercropping soil with *C. indica* showed lower species abundance in comparison with soil without *C. indica* (ACE index 112 and 125,6 respectively). In contrast, soil with *C. indica* has more taxa richness in comparison with soil without *C. indica* (Fisher index 278 and 230,4, respectively)..

Keywords: 16S rRNA, metagenomic analysis, Rhizosphere bacteria, Intercropping system, *Canna indica*

1. Introduction

Rubber tree (*Hevea brasiliensis*) is one of economically important crop plants in some countries, especially in developing countries. With about 3.5 million hectare rubber tree plantation and total production about 2.7 tones raw rubber (Statistik Perkebunan, 2010), Indonesia is one of a major rubber producer in the world. Mostly rubber tree plantations in Indonesia are managed by traditional farmers (84.66%) and the rest by private corporates and Government (BPS, 2009). However, for the last couple years world rubber price is showing a declining trend. It makes rubber plantation is not any more interesting for most of traditional farmers. Increasing production cost especially fertilizer and labor costs that is not followed by an increase in rubber price is major problem in rubber tree



production. As the results, the agronomy and management system of rubber plantations belonged traditional farmers and Indonesian government are no longer ideal. Thus, low productivity and quality of raw rubber are the consequences faced by raw rubber production.

Some efforts have been proposed to resolving this problems, one of them is using intercropping system rubber tree with other suitable plants. Intercropping system is a practice of growing a crop with other crops in close proximity in the same growing season. Basically, intercropping system is application of ecological principle and an efficiency effort of natural resources usage (Duan et al, 2011).

On the rubber plantation, usage of arable land under rubber tree canopy is a challenging idea. Since the rubber canopy is quite dense allowing little solar radiation through understory. It has been reported only about 20% of incoming solar radiation is available under 4-5 years old rubber canopy (Ibrahim, 1991), thus determining of properly plants for intercropping system with rubber tree is critical factor for success intercropping system. Indeed, intercropping system has been shown could enhance soil nutrient contents such as N and P, soil enzyme activities and crop yield (Zhou et al, 2011; Ghosh et al, 2006).

Another important factor which appears as the result of intercropping system is change of soil microbial community. Several studies showed that structure of microbial community and metabolic capabilities could reveal soil quality conditions (Yao et al, 2006). Microbial community has important role in determining of the soil ecosystem and the conversion of soil nutrient cycling. Nitrogen, sulfur and phosphorus cycles, as well as ecosystem functioning are some crucial role of soil microbes. Later on it could change soil structure formation, organic matter decomposition, and toxin removal (Acosta-Martinez *et al.*, 2010; Karlen *et al.*, 1997; Gomes *et al.*, 2003). Indeed, soil microbial diversity is important for maintaining good quality of agricultural soil (Acosta-Martinez *et al.*, 2010; Garbeva *et al.*, 2004; Janvier *et al.*, 2007).

Since soil bacteria is huge not only in diversity but also in abundance, and most of bacteria are unculturable (Nihorimbere et al. 2011), thus using of standard culturing method will not enough to describe the richness of soil bacteria. Using high throughput sequencing technique (Next Generation Sequencing/NGS) of specific conserved genes or genomic fragments such as 16S rRNA, this information become accessible (Riesenfeld et al. 2004; Effendi et al, 2019; Ghosh et al. 2019). Using metagenomic analysis of 16S rRNA gene of soil microbial that isolated around rhizosphere area of *C. indica* root system, this research was aimed to investigate impact of intercropping of *C. indica* in the rubber plantation on diversity and richness of soil microbial.

2. Materials and Methods

2.1. Sampling

Soil samples were collected from rubber tree plantation of PTPN VIII in Subang, West Java, Indonesia. Soil samples consisted of 500 g soil samples were collected from three different sites nearby rhizosphere areas of Rubber tree-*Canna* plants (G) intercropping area or from three different sites nearby rhizosphere areas of *Rubber* trees without *Canna* plants nearby. Each three soil samples were homogenized and taken 50 g for DNA isolation. This procedure was repeated twice for each point locations. The soil samples were labeled according to location and kept in 4 °C until used.

2.2. DNA extraction and molecular works

Whole genomic DNA was extracted using PowerSoil DNA kit (MoBio). DNA extraction was done following manual procedure of the kit. A 0.25 gr soil of each sample were used as source of the

genomic DNA. All procedures were performed aseptically for avoiding contamination. The DNA results were quantified using NanoDrop and checked its quality using gel electrophoresis. The DNA was used as a template for PCR. A 2 μ L of DNA was added to 10 μ L PCR mix (GoTaq[®] Green Master mix-PROMEGA) and 1 nmol of each forward and reverse 16S rRNA primer. Reaction was performed 35 cycles which consisted of 30 sec at 94°C and continued with 57°C of annealing for 20 sec, followed with 2 min 72°C for elongation. A region V4 of 16S rRNA gene was amplified with primer F515 (5'-GTGCCAGCMGCCGCGGTAA-3') and 907R (5'-CCGTCAATTCMTTTRAGTTT-3') (Lane 1991). The PCR products were purified and subjected for automated Illumina Miseq platform (1st BASE-Malaysia) after the PCR products were normalized in equimolar amounts.

2.3. Sequences analysis

Raw sequence data generated from Illumina Miseq platform were processed in QIIME Ver 6.0 (Caporaso et al. 2010). All sequences shorter than 150 bp or longer than 600 bp were removed from downstream processing. Read were then aligned with 16S rRNA SILVA database (www.arb-silva.de) and GRD database (metasystems.riken.jp/grd/), then followed inspected for chimeric errors. "Species-level" of OTU was used in analysis, thus reads then were clustered at 97% similarity into OTUs. In this step, rare OTUs with only one (singleton) or two reads (doubleton) were deleted from downstream processing. Taxonomic assignment was carried out with the RDP Classifier (Wang et al. 2007).

2.4. Data analysis

Alpha diversity analysis was conducted using qiime2 for evenness and Faith phylogenetic distance index, and phyloseq for Shannon and Chao1 diversity index. Beta diversity was analyzed with Bray-Curtis and Weighted Unifrac index in Qiime2 Alpha and Beta diversity analysis were performed using Explicit ver 2.10.5 software (Robertson et al. 2013). Statistical comparison of alpha diversity between samples was carried out with Excel (Microsoft) whereas statistically different (*P-value*) was calculated with T-test. Sample-specific OTUs (showing significantly different relative abundances between samples) were assessed by T-test. Heat map which showed relative abundances between samples in certain taxa level was generated with Explicit ver 2.10.5 software.

3. RESULT AND DISCUSSION

After filtering total reads based on quality control of raw read data, total of 764,951 sequences were collected from 4 soil samples. About 300,570 singleton and chimera were identified and removed. Altogether 25,243 OTUs were identified after were aligned with SILVA database and GRD database with similar proportion among the 4 soil samples (Table 1).

Table 1. Number of reads after filtering and OTUs

Samples	Raw reads	Chimera and singleton	Number of OTU
With Canna (G20)-intercropping	177219	65846	6254
With Canna (G40)-intercropping	229973	92792	6546
Without Canna (TG20)	165555	62485	5810
Without Canna (TG40)	192204	79447	6633

3.1. Diversity and Structure of bacterial community

Analysis of microbial Diversity based on Shannon-Winner (H') index showed that the highest index value was occurred in the TG40 sample ($H' = 3.0506$) and the lowest value at G20 ($H' = 2.9467$) (Table 2). According to Fanani *et al.* (2013) a low H' values ($H' \leq 1$) indicate a low microbial diversity, whereas sample with $H' \geq 3$ indicate a high diversity in microbial community. Our data indicated all

soil samples have high microbial diversity, except soil sample from intercropping area collected from 20 cm soil depth which has moderate microbial diversity.

Table 2. Shannon-Wiener (H') and Simpson (H_2) diversity index values on G and TG at each soil depth conditions.

Sample	Indeks Shannon-Wiener (H')	Indeks Simpson (H_2)
With Canna - intercropping (G20)	2.9467	0.093
With Canna - intercropping (G40)	3.0236	0.085
Without Canna (TG20)	3.0148	0.085
Without Canna (TG40)	3.0506	0.078

The lowest H' index value for the G20 can be expected due to the presence of certain types of microbes that dominate the G20 area, so that microbial diversity has decreased. According to Cover (1991), the value of the diversity index will be maximized when all individuals are equally represented. The higher diversity value shows that a community has a relatively even number of individuals and no species domination. Microbial diversity is influenced by the physical chemistry of soil (Meilani et al, 2012; Effendi et al, 2019). Hill *et al.* (2003) reported that diversification function of a land will experience a decline, especially in acacia forests, oil palm plantations and rubber plantations.

Moreover, the Simpson index (H_2) at the depth of the G20 has a higher index value of ($H_2=0.093$) while the lowest value is found in the TG40 sample ($H_2=0.078$). Based on the analysis results, the index value is less than 0.5 which means that each class at the four individual depths does not dominate. This condition shows that the microbial and ecological communities are still in a stable state.




Figure 1. Relative abundance of the phyla from 4 soil samples (G20, G40, TG20, TG40).

Based on the number of observed OTUs, 97% OTUs were occupied by bacteria and the rest was Archaea (3%). Figure 1 showed relative abundance of phyla composition that identified. Although the 4 soil samples showed similar relative abundance, it is clear that some phyla abundance showed obvious variations between different soil conditions. A total 34 phyla were affiliated to bacteria phyla in database, whereas 2 remaining groups were still unclassified. The most abundant phyla of *Acidobacteria*, *Chloflexi* and *AD3* were found in all soil samples with relative similar proportion (25.8%, 21.6%, 16.7% respectively in sample G dan 23.5%, 20.5%, 16.2% respectively in samples TG). The next 6 phyla (*Verrucomicrobia*, *Proteobacteria*, *Planctomycetes*, *GLA15*, *Nitrospirae* and *Actinobacteria*) were found in all soil samples at a relative abundance that higher than 2% but lower than 8% with obvious variations. The relative abundance of *Verrucomicrobia* were relative similar in G20, TG20 and TG40, but lowest in G20. *Proteobacteria* were found more abundant in TG samples than in G samples.

Most of the phyla have similar abundance in the soil samples, but some phyla were identified different in abundance. The phyla of BHI80.139 ($p < 0.00$), *Proteobacteria* ($p < 0.02$), *Cyanobacteria* ($p < 0.03$), TM6 ($P < 0.03$), and WS3 ($p < 0.02$) were found significantly different in their abundances. The complete data is shown in table 3. *Proteobacteria* and *Cyanobacteria* were found more abundant in TG samples. Both phyla were known involving in biogeochemical cycle particularly in nitrogen fixation (Vijayan and Ray, 2015). The dominant phylum of *Proteobacteria* was also reported in some publications in various soil types (Sul et al, 2013, Chu et al, 2010; Nacke et al, 2011). The bacterial community composition and diversity were largely affected by environment factors, soil pH and plant species (Wei et al, 2017). The soil pH was the key factor influencing bacterial diversity, with lower pH associated with less diverse communities (Wei et al, 2017). However, in this study, soil conditions and pH were similar. Soil analysis of TG and G samples indicated pH of 4.3 and 4.6, respectively. Moreover, their C organic contents were also almost same 1.54% and 1.95%, respectively. It indicates that the presence of Canna plant may contribute significant in abundance of *Proteobacteria* and *Cyanobacteria*.

Vijayan and Ray (2015) indicated that there is positive correlation on the total number of *Cyanobacteria* inhabiting tropical paddy wetland to total nitrogen in these soils. However, our data showed that the N total in the TG and G samples has no significant differences (0.45% and 0.36% respectively). According to soil analysis data, both soils conditions indicated low fertile soil condition. Table 3. Signification of abundance of the phyla group between Canna samples (Cs) and without Canna samples (TGs)

Phylum	<i>Canna indica</i> (G)	Without <i>C. indica</i> (TG)	<i>p</i> -value
<i>AD3</i>	0.33397 ± 0.03336	0.32471 ± 0.02294	0.46
<i>Acidobacteria</i>	0.51663 ± 0.01456	0.47099 ± 0.01643	0.20
<i>Actinobacteria</i>	0.06958 ± 0.00217	0.09389 ± 0.00446	0.07
<i>Armatimonadetes</i>	0.01092 ± 0.00145	0.01410 ± 0.00092	0.23
<i>BHI80.139</i>	0.00003 ± 0.00002	0.00000 ± 0.00000	0.00
<i>BRC1</i>	0.00006 ± 0.00000	0.00000 ± 0.00000	0.09
<i>Bacteroidetes</i>	0.00342 ± 0.00081	0.00392 ± 0.00078	0.42
<i>Chlamydiae</i>	0.00139 ± 0.00023	0.00159 ± 0.00017	0.38
<i>Chlorobi</i>	0.00018 ± 0.00002	0.00010 ± 0.00001	0.09
<i>Chloroflexi</i>	0.43239 ± 0.00326	0.41125 ± 0.01023	0.21
<i>Cyanobacteria</i>	0.00109 ± 0.00006	0.00169 ± 0.00005	0.03

<i>Elusimicrobia</i>	0.00227 ±0.00022	0.00316 ±0.00032	0.18
<i>FCPU426</i>	0.00283 ±0.00012	0.00311 ±0.00017	0.28
<i>Fibrobacteres</i>	0.00002 ±0.00001	0.00003 ±0.00002	0.41
<i>Firmicutes</i>	0.00069 ±0.00003	0.00125 ±0.00019	0.15
<i>GAL15</i>	0.10245 ±0.00512	0.07792 ±0.00306	0.09
<i>Gemmatimonadetes</i>	0.01192 ±0.00116	0.01736 ±0.00005	0.07
<i>NC10</i>	0.00148 ±0.00025	0.00052 ±0.00013	0.12
<i>Nitrospirae</i>	0.06793 ±0.00064	0.04387 ±0.00494	0.06
<i>OD1</i>	0.00286 ±0.00012	0.00392 ±0.00055	0.22
<i>OP11</i>	0.00003 ±0.00002	0.00000 ±0.00000	0.21
<i>OP3</i>	0.00005 ±0.00003	0.00003 ±0.00002	0.38
<i>Planctomycetes</i>	0.08480 ±0.00910	0.12221 ±0.00413	0.10
<i>Proteobacteria</i>	0.11121 ±0.00074	0.14108 ±0.00336	0.02
<i>Synergistetes</i>	0.00000 ±0.00000	0.00004 ±0.00002	0.21
<i>TM6</i>	0.00034 ±0.00005	0.00029 ±0.00008	0.03
<i>TM7</i>	0.00301 ±0.00032	0.00254 ±0.00010	0.28
<i>Verrucomicrobia</i>	0.09866 ±0.01891	0.13636 ±0.00072	0.21
<i>WPS.2</i>	0.01723 ±0.00170	0.01625 ±0.00148	0.42
<i>WS3</i>	0.00294 ±0.00033	0.00679 ±0.00022	0.02

CONCLUSION

This study showed that intercropping of *Hevea brasiliensis* with *Canna indica* may contribute in the status of diversity and abundance of soil bacteria. Most of the phyla members were not changed significantly after intercropping of *Canna indica*. However, *Proteobacteria* and *Cyanobacteria* were found lower abundance in the soil without *Canna indica* (TG samples), whereas both phylum has been reported positively correlated with biogeochemical cycles in the soil particularly in determining of nitrogen content in the soil.

ACKNOWLEDGEMENTS

This research was supported financially by research grant of PDUPT 2018 from The Ministry of Research Technology and Higher Education-Republic of Indonesia and technically supported by PTPN VIII West Java Indonesia.

Reference

- [1] Acosta-Martínez, V., G. Burow, T.M. Zobeck and V.G. Allen. 2010. Soil microbial communities and function in alternative systems to continuous cotton for the Texas high plains. *Soil Sci. Soc. Am. J.*, 74: 1181-1192.
- [2] Badan Pusat Statistik 2009. Ekspor-Impor Karet. Badan Pusat Statistik Jakarta.
- [3] Cover TM, Thomas JA. 1991. *Elements of Information Theory*. New York: John Wiley and Sons Inc.
- [4] Chu H, Fierer N, Lauber CL, Caporaso JG, Knight R, Grogan P. 2010. Soil bacterial diversity in the Arctic is not fundamentally different from that found in other biomes. *Environ Microbiol.* 12(11):2998-3006.
- [5] Direktorat Jenderal Perkebunan. 2010. *Statistik Perkebunan Indonesia 2009-2011*. Komoditas Karet, Direktorat Jenderal Perkebunan, Jakarta 2010.
- [6] Duan, F., Wang, X.Y., Gao, Z.H. 2011. Causes and control measures of continuous cropping obstacle in horticultural crops. *Acta Agriculturae Jiangxi*, 23(3): 34-39

- [7] Effendi Y, Pambudi S, Pancoro A. 2019. Metagenomic analysis of *Fusarium oxysporum* f.sp. *cubense*-infected soil in banana plantation, Sukabumi, Indonesia. *Biodiversitas* 27 (7) 1939-1945
- [8] Fanani A, Rohman F, Sulasmi, Eko S. 2013. Karakteristik Komunitas Herba di Hutan Jati Resort Pemangkuan Hutan (RPH) Dander Petak 12 B Kabupaten Bojonegoro. Malang: Universitas Negeri Malang.
- [9] Garbeva, P., J.A. van Veen and J.D. van Elsas. 2004. Microbial diversity in soil: selection of microbial populations by plant and soil type and implications for disease suppressiveness, *Annu. Rev. Phytopathol.*, 42: 243-270.
- [10] Ghosh, P.K., Mannab, M.C., Bandyopadhyayb, K.K., et al. 2006. Interspecific interaction and nutrient use in soybean/sorghum intercropping system. *Agron J*, 98: 1097-1108.
- [11] Ghosh A, Mehta A, Khan AM. 2019. Metagenomic Analysis and its Applications. *Encyclopedia Bioinformatics Comput Biol* 3: 184-193
- [12] Gomes, N.C., O. Fagbola, R. Costa, N.G. Rumjanek, A. Buchner, L. Mendona-Hagler and K. Smalla. 2003. Dynamics of fungal communities in bulk and maize rhizosphere soil in the tropics. *Appl. Environ. Microbiol.*, 69: 3758-3766.
- [13] Hill TCJ, Walsh KA, Harris JA, Molett BF. 2003. Using Ecological Diversity measures with Bacterial Communities. London: University of East London.
- [14] Ibrahim A.G., 1991. Influence of rubber canopy on intercrop productivity. *Trans.MalaysianSoc.Pl. Physiol.*2,75-'79.
- [15] Janvier, C., F. Villeneuve, C. Alabouvette, V. Edel-Hermann, T. Maitelle and C. Steinberg. 2007. Soil health through soil disease suppression: which strategy from descriptors to indicators. *Soil Biol. Biochem.*, 39: 1-23.
- [16] Karlen, D.L., M.J. Mausbach, J.W. Doran, R.G. Cline, R.F. Harris and G.E. Schuman. 1997. Soil quality: a concept, definition, and framework for evaluation. *Soil Sci. Soc. Am.J.*, 61: 4-10
- [17] Munthe VY, Aryawati R, Isnaini. 2012. Struktur Komunitas dan Sebaran Fitoplankton di Perairan Sungang Sumatera Selatan. *J Maspari*. 4:122-130
- [18] Nacke H, Thürmer A, Wollherr A, Will C, Hodac L, Herold N, Schöning I, Schrupf M, Daniel R. 2011. Pyrosequencing-based assessment of bacterial community structure along different management types in German forest and grassland soils. *PLoS One*. 6(2):e17000.
- [19] Nihorimbere V, Ongena M, Smargiassi M, Thonart P. 2011. Beneficial effect of the rhizosphere microbial community for plant growth and health. *Biotechnol Agron Soc Environ* 15: 327-337.
- [20] Riesenfeld CS., Schloss PD, Handelsman J. 2004. METAGENOMICS: Genomic Analysis of Microbial Communities. *Annu Rev Genet* 38: 525-52
- [21] Robertson CE, Harris JK, Wagner BD, Granger D, Browne K, Tatem B, Feazel LM, Park K, Pace NR, Frank DN. 2013. Explicet: graphical user interface software for metadata-driven management, analysis, and visualization of microbiome data. *Bioinformatics* 29 (23): 3100-3101
- [22] Saadatnia H, Riahi H. Cyanobacteria from paddy fields in Iran as biofertilizer. *Plant Soil*. 2009;55:207–212
- [23] Sul WJ, Asuming-Brempong S, Wang Q, Tourlousse DM, Penton CR, Deng Y, et al. 2013. Tropical agricultural land management influences on soil microbial communities through its effect on soil organic carbon. *Soil Biol Biochem*. 65:33–8
- [24] Sutrisna T, Umar MR, Suhadiyah S, Santosa S. 2018. Keanekaragaman dan Komposisi Vegetasi Pohon pada Kawasan Air Terjun Takapala dan Lanna di Kabupaten Gowa, Sulawesi Selatan. *J Biologi Makassar*. 3:12-18.
- [25] Vijayan D, Ray JG. 2015. Ecology and diversity of Cyanobacteria in Kuttanadu paddy wetlands, Kerala, India. *Am J Plant Sci*. 6:2924–2938.
- [26] Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naïve bayesianclassifier for rapid assignment

- of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 73: 5261-5267
- [28] Wei Z, Hu X, Li X, Zhang Y, Jiang L, Li J, et al. 2017. The rhizospheric microbial community structure and diversity of deciduous and evergreen forests in Taihu Lake area, China. *PLoS ONE* 12(4): e0174411
- [29] Yao, H.Y., Daniel, B., Wei, S. 2006. Soil microbial community structure and diversity in a turfgrass chronosequence: Land-use change versus turfgrass management. *Appl Soil Ecol*, 34(2): 209-218
- [30] Zhao, Q.L., S.X. Liang, S.X. Z.M. Ling and L.Z. Tai. 2011. Studies on Soil microorganism quantities and soil enzymes activities in the garlic-cotton and wheat-cotton intercropping systems. *Plant Nutr. Fert. Sci.*, 17: 1474-1480.