

Klarifikasi: (lampiran bukti kehadiran)

Poster yang berjudul: “Auxin-Binding-Protein1 (ABP1), the Second Auxin Receptor“ dipublikasikan dalam bentuk poster presentation pada 20th International Conference on Plant Growth Substance (20th IPGSA) Tarragona, Spain, 28th Juni – 2nd July 2010 yang diselenggarakan Unversiti Rovira I Virgili. Penulis sebagai first author (penulis utama). Konferensi ini tidak menyediakan adanya prosiding sehingga publikasi ini **tergolong tidak dimuat dalam prosiding**. Informasi tentang konferensi (paniti penyelenggara, peserta dan poster/makalah yang dipresentasikan) terinfokan pada book of abstract (dilampirkan). Hanya peserta yang mempresentasikan makalahnya secara oral (Oral presentation) yang diminta untuk menyerahkan artikelnya, sedangkan publikasi berupa presentasi poster, tidak disyaratkan adanya artikel. Sebagai bukti keikutsertaan kami dilampirkan

- Sertifikat kehadiran
- Daftar committe of conference
- Cover book of abstract dan daftar kegiatan
- Screenshot nomer poster yang dipresentasikan
- File yang berisi URL informasi tentang 20th IPGSA (book of programme)

Auxin-binding protein 1 (ABP1), the second auxin receptor

Yunus Effendi¹, Steffen Rietz², Urs Fischer³, Günther F.E. Scherer¹

1) Molecular Developmental Physiology, University Hannover, Herrenhäuser Str. 2, D-30419 Hannover, Germany;
2) Max Planck Institute for Plant Breeding Research, Cologne; 3) Forstbotanik und Baumphysiologie University Göttingen.
Scherer@zier.uni-hannover.de

Leibniz
Universität
Hannover

Abstract

Despite of knowing the 3-dimensional structure ABP1 is not fully acknowledged as an auxin receptor. We used the homozygous lethal ABP1 insertional mutant (Chen et al. 2001) which is viable in the heterozygous *abp1/ABP1* state. *abp1/ABP1* seedlings are defect in phototropism and gravitropism of roots and shoots. Those populations are composed of a major slow reacting and a minor normal reacting group. *abp1/ABP1* seedlings show strong root slanting, longer hypocotyls, and only slightly increased lateral root numbers. Root auxin responses (lateral roots, main root length) in *abp1/ABP1* seedlings are only slightly less sensitive than in wt. In short and long days *abp1/ABP1* plants flower earlier. They have more branches and decreased main stem diameter, indicating decreased apical dominance. Auxin-induced genes (qPCR of: *IAA2*, *IAA11*, *IAA12*, *IAA13*, *IAA14*, *IAA19*, *IAA20*, *SAUR9*, *SAUR15*, *SAUR23*, *GH3.5*, *ABP1*) respond to auxin (0.1 μM/1 μM/10 μM) 2-10-fold stronger in wt than in *abp1/ABP1* seedlings (30 & 60 min). Auxin content and uptake of auxin in *abp1/ABP1* seedlings is not distinguishable from wt. Basipetal auxin transport in *abp1/ABP1* roots is slower than in wt. Thus ABP1 is a receptor with probable functions in auxin transport and gene regulation. The necessary functional link to TIR1-linked gene regulation could be provided by phospholipase(s) A (FEBS Lett. 2007, 581:4205-4211).

Defects in auxin-related phenotypes in heterozygous *abp1/ABP1* plants





Fig. 1. Early flowering phenotype of wild type Ws and heterozygous *abp1/ABP1* plants grown in short days (8/16 light/day) (A, B, C) or long days (16/8 light/day) (D, E). (A) In short days grown progeny of seeds from a kanamycin-resistant heterozygous *abp1/ABP1* plant (in background) and Ws wild type plants (front row). Plants were grown in short days and ordered, those with open flowers to the back and with labeled small white tags, and those not having flowers, with small red tags. (B) On short day and (D) long day, *abp1/ABP1* produced flower earlier [(p<0.01) and (p<0.05) respectively]. WT has more rosette leaves (p<0.05) on Long day (C) and also short day (p<0.05)(D). Flowering was recorded as first open flower. Rosette leaves and cauline leaves were counted at the time of bolting. Cauline leaf numbers were not different in all cases. All plants were PCR-genotyped.

Defect in apical dominant of heterozygous *abp1/ABP1* plants


Fig. 3. Apical dominance of wild type Ws and heterozygous *abp1/ABP1* plants grown in long days (16/8h light/dark) or short days (8h/16h light/dark). (A) Long day wild type Ws plant. (B) Long day heterozygous *abp1/ABP1* plant. Red arrow: main stem; yellow arrows: lateral stems. Plant genotypes were determined by PCR. Note the lower number of rosette leaves and absence of prominent main stem in mutant plants. One out of three experiments is shown. (C) Stem thickness and stem number of wild type Ws long day plants (n as indicated; S.E.). (D) Stem thickness and total number of stems of heterozygous *abp1/ABP1* plants (n as indicated; S.E.). (E) Stem thickness and stem number of wild type short day Ws plants (n as indicated; S.E.). (F) Stem thickness and total number of stems of short day heterozygous *abp1/ABP1* plants (n as indicated; S.E.). Plant genotypes were determined by PCR. Numbers on top of bars: total number of branches for this branch class. One out of two experiments is shown.

Fig. 3




Regulation of early auxin-induced genes in heterozygous *abp1/ABP1* showed slightly defect in comparison to WT

Fig. 7



Auxin sensitivity of *abp1/ABP1* seedling

Fig. 2. Auxin sensitivity of *abp1/ABP1* seedlings. All seedlings were pre-grown on vertical agar plates without auxin for 4 d, and then transferred to plates containing increasing concentrations of IAA. The *abp1/ABP1* seedlings were selected as having a strong slanting angle from the segregating population after 3 d and both Ws and mutant seedlings were transferred to a fresh plate for 4 d. Response to auxin of (A) relative main root length, (B) lateral root number (C) relative hypocotyl length. (n=20 wild type Ws; n= 20 *abp1/ABP1*; S.E.).



Phenotypic appearance and responses to gravity and light




Fig. 4

Fig. 4. Phenotypic appearance and responses to gravity and light in heterozygous *abp1/ABP1* seedlings (A). Phenotype of 2 weeks old WT and heterozygous *abp1/ABP1* seedlings. (B) Heterozygous *abp1/ABP1* produce greater slanting angles (n=20; S.E., p<0.01) and longer hypocotyl (n=20; S.E., P<0.01) (C). (D) Representative images of 4 d old seedlings after 24 h gravitropic response and (E) their graphic quantification of hypocotyl curvature angle (n=180; S.E.). (F) Representative images of 7 d old seedlings after 24 h gravitropic response and (G) their graphic quantification of root curvature angle (n=96; S.E.). (H) Representative images of 4 d old seedlings after 10 h phototropic response and (I) their graphic quantification (n=80; S.E.). Open diamonds: seeds from a kanamycin-resistant *abp1/ABP1* plant; Filled squares: Ws

No different in auxin uptake between heterozygous *abp1/ABP1* and Wild type, but auxin moved slower in heterozygous *abp1/ABP1* plants in basipetal direction

Fig. 5




Fig. 5. Auxin uptake into wild type and *abp1/ABP1* seedlings. Plants were grown, selected, and treated with auxin as described in the legend for Fig. 7. Analysis of IAA content was performed with quadrupole GC-MS system (Agilent) and results from three experiments were pooled. (A) Treatment with 0.1 μM IAA. (B) Treatment with 1 μM IAA. (C) Treatment with 10 μM IAA. Wild type plants: filled squares, *abp1/ABP1*: open diamonds (n= 4-6; S.D.).




Fig. 6

Fig. 6. Polar auxin transport in wild type and *abp1/ABP1* roots. (A, B) Acropetal transport in the presence of 15 μM NPA (A) or without (B). Two consecutive 5 mm pieces near the source and the residual third root piece were counted as indicated on the left side of the figure. No statistically significant differences in any segment. (C, D) No significant different in basipetal transport in the presence of 15 μM NPA (C) but there were statistically significant differences (p<0.05 one asterisk (*) in middle segment and at <0.001 two asterixes (** in lower segment) in basipetal transport without NPA (D)

Fig. 7 Regulation of early auxin-regulated genes and of ABP1 in light-grown wild type (Ws) and *abp1/ABP1* seedlings. 14 days old seedlings were grown in 1/2 MS agar and treated with 0.1 μM, 1 μM, or 10 μM IAA for 30 min and 60 min. Each data were collected from 2 (B) or 3 (A, C) biological replications and 3 technical replications independently. Twelve genes (*IAA2*, *IAA11*, *IAA12*, *IAA13*, *IAA14*, *IAA19*, *IAA20*, *SAUR9*, *SAUR15*, *SAUR23*, *GH3.5*, *ABP1*) were investigated using quantitative Real Time PCR method (qRT-PCR) and the results showed that those genes are expressed 2-10-fold stronger in wt than in *abp1/ABP1*. (A) Treatment with 0.1 μM IAA. (B) 1 μM IAA. (C) 10 μM IAA. Wild type: open squares. *abp1/ABP1*: open diamonds.

Conclusion: ABP1 is a powerful receptor which regulates genes, likely with a functional link to TIR1. The heterozygous *abp1/ABP1* mutant is defect in responses requiring polar auxin transport.

References:

Scherer, G.F.E., Zahn, M., Callis, J. and Jones, A.M. (2007) A role for phospholipase A in auxin-regulated gene expression. FEBS Lett. 581, 4205-4211



IPGSA CONFERENCE 2010

20th International Conference
on Plant Growth Substances

28th June to 2nd July 2010
Universitat Rovira i Virgili
Tarragona - Spain

www.ipgsa.org

Organizers:

Montserrat Pagès (Barcelona)
Lluís Arola (Tarragona)
Juan Carbonell (Valencia)

Plenary Speakers:

Moto Ashikari (Japan)
Bonnie Bartel (USA)
Jiri Friml (Belgium)
Markus Grebe (Sweden)
Veronica Grieneisen (UK)
Erwin Grill (Germany)
Luis Herrera-Estrella (Mexico)
Tetsuya Higashiyama (Japan)
Joe Kieber (USA)
Ottoline Leyser (UK)
Eliezer Lifschitz (Israel)
Salomé Prat (Spain)
Paul Staswick (USA)
Zhiyong Wang (USA)
Shinjiro Yamaguchi (Japan)



	June 28 (Monday)	June 29 (Tuesday)	June 30 (Wednesday)	July 1 (Thursday)	July (Friday)		
09:00	Registration (opens every day)	P2 Hormone Biosynthesis and Transport <i>Shinjiro Yamaguchi (Japan)</i>	P3 Hormone Perception and Signaling <i>Joe Kieber (USA)</i>	PS09 Evolution of Plant Hormone Signaling PS10 Root Development PS11 Abiotic Stress PS12 Chemical Biology	P4 Hormones and Environment <i>Salomé Prat (Spain)</i> <i>Moto Ashikari (Japan)</i>		
09:45		<i>Paul Staswick (USA)</i>	<i>Bonnie Bartel (USA)</i>				
10:30		Coffee break	Coffee break	Coffee break	Coffee break		
11:00		<i>Tetsuya Higashiyama (Japan)</i>	<i>Zhiyong Wang (USA)</i>	PS09 Evolution of Plant Hormone Signaling PS10 Root Development PS11 Abiotic Stress PS12 Chemical Biology	Steve Penfield (UK)		
11:30		Opening					
11:45	Silver Medal Lectures <i>Joanne Chory (USA)</i>	<i>Jiri Friml (Belgium)</i>	<i>Erwin Grill (Germany)</i>		<i>Luis Herrera-Estrella (Mexico)</i>		
12:30	<i>Tokao Yolota (Japan)</i>	Lunch & Posters	Lunch & Posters	Lunch & Posters	Closing		
13:00							Farewell Lunch
13:15	<i>Mark Estelle (USA)</i>						
14:00	Lunch						
15:00	P1 Hormones and Development <i>Ottoline Leyser (UK)</i>	PS01 Hormone Biosynthesis PS02 Seed Development and Germination PS03 Auxin Transport PS04 Light Responses	PS05 Hormone Perception and Signaling PS06 Shoot Development PS07 Defense responses PS08 Systems Biology	PS13 Hormone Interactions PS14 Reproductive Development PS15 Hormones and Biotechnology PS16 Epigenetics/Small RNAs			
15:45	<i>Eliezer Lifschitz (Israel)</i>						
16:30	Coffee break	Coffee break	Coffee break	Coffee break			
17:00	<i>Markus Grebe (Sweden)</i>	PS01 Hormone Biosynthesis PS02 Seed Development and Germination PS03 Auxin Transport PS04 Light Responses	PS05 Hormone Perception and Signaling PS06 Shoot Development PS07 Defense responses PS08 Systems Biology	PS13 Hormone Interactions PS14 Reproductive Development PS15 Hormones and Biotechnology PS16 Epigenetics/Small RNAs			
17:45	<i>Veronica Grieneisen (UK)</i>						
18:30	Wellcome Reception	Tarragona Roman Ruins visit		IPGSA Business Meeting			
19:30							
20:00				congress Dinner			

IPGSA CONFERENCE 2010

20th International Conference
on Plant Growth Substances

28th June to 2nd July 2010
Tarragona (SPAIN)

I309

CERTIFICATE OF ATTENDANCE

We hereby certify that,


MR. YUNUS EFFENDI

has attended the


20th International Conference on Plant Growth Substances

***held at the Universitat Rovira i Virgili, Tarragona (Spain), on 28th
June to 2 July 2010.***

**Organizing Secretariat
Grupo Geyseco, S.L.**



IPGSA
CONFERENCE
2010
28th June to 2nd July 2010
Universitat Rovira i Virgili and
Palau de Congressos
TARRAGONA - SPAIN
20th International Conference
on Plant Growth Substances



User: Password: >>

- Home
- Welcome
- Committees**
- Sessions and Chairs
- Program at a glance
- Scientific Program
- General Information
- Registration & Accommodation
- Poster Sessions
- Tarragona
- Sponsors
- Downloads
- Personal Area

Committee

Organizers:

- Montserrat Pagès (Barcelona), chair
- Lluís Arola (Tarragona)
- Juan Carbonell (Valencia)

Local Organizing Committee

- Adela Goday (Barcelona)
- Pere Puigdomenech (Barcelona)
- Antonio Tiburcio (Barcelona)
- Francisco Culiáñez (Valencia)
- José Luis García-Martínez (Valencia)
- Isabel López-Díaz (Valencia)

Scientific Committee

- Tadao Asami (Japan)
- Bonnie Bartel (US)
- Caren Chang (US)

- PS05-18** **STRUCTURAL BASIS FOR GIBBERELLIN RECOGNITION BY ITS RECEPTOR GID1**
Miyako Ueguchi-Tanaka (Japan)
- PS05-19** **MIS-REGULATION OF EARLY AUXIN-INDUCED GENES IN PHOSPHOLIPASE A KNOCKOUTS**
Corinna Labusch (Germany)
- PS05-20** **CHEMICAL SCREENING AND CHARACTERIZATION OF INHIBITORS FOR GIBBERELLIN RECEPTOR**
Jung-min Yoon (Japan)
- PS05-21** **A POSSIBLE CYTOSOLIC ACTIVITY FOR THE GA RECEPTOR, GID1**
Sivan Sagiv (Israel)
- PS05-22** **DOMAIN REQUIREMENTS FOR DIFFERENT ROLES OF THE ETR1 ETHYLENE RECEPTOR**
Brad Binder (USA)
- PS05-23** **TRANSCRIPT PROFILING OF CYTOKININ ACTION IN ARABIDOPSIS ROOTS AND SHOOTS**
Brenner Wolfram (Germany)
- PS05-24** **THE IMA GENE ENCODES AN ADAPTER MINI ZINC FINGER PROTEIN REGULATING THE DEVELOPMENT OF FLORAL MERISTEM IN TOMATO.**
LEBLOND JULIE (FRANCE)
- PS05-25** **QUANTITATIVE DIFFERENTIAL ANALYSIS OF PLANT PROTEOME IN A RESPONSE TO HORMONAL REGULATIONS**
Marketa Zdarska (Czech Republic)
- PS05-26** **DYNAMIC STRUCTURE AND BINDING SPECIFICITY OF THE RECEIVER DOMAIN OF SENSOR HISTIDINE KINASE CKI1**
Tomas Klumpler (Czech Republic)
- PS05-27** **ATPLA-I, A PHOSPHOLIPASE A WITH FUNCTIONS IN GRAVITROPISM, PHOTOTROPISM AND ROOT TIP MOVEMENTS**
Günther Scherer F.E. (Germany)
- PS05-28** **AUXIN-BINDING PROTEIN1 (ABP1), THE SECOND AUXIN RECEPTOR**
Yunus Effendi (Germany)
- PS05-29** **MUTATION OF TRNA SPECIFIC ISOPENTENYLTRANSFERASES AFFECTS ARABIDOPSIS PHENOTYPE**
Silvia Gajdosova (Czech Republic)