

[Ad covered content](#)[Ad was inappropriate](#)[Seen this ad multiple times](#)[Not interested in this ad](#)

Journal of Applied Phycology

COUNTRY[Netherlands](#)

Universities and research institutions in Netherlands

SUBJECT AREA AND CATEGORY[Agricultural and Biological Sciences](#)
[Aquatic Science](#)
[Plant Science](#)[Ad covered content](#)[Ad was inappropriate](#)[Seen this ad multiple times](#)[Not interested in this ad](#)**PUBLISHER**[Springer Netherlands](#)**H-INDEX**

101

PUBLICATION TYPE[Journals](#)**ISSN**

15735176, 09218971

COVERAGE


1989-2020

INFORMATION[Homepage](#)[How to publish in this](#)


FIND SIMILAR JOURNALS ?

1 Algal Research NLD 55% similarity	2 Botanica Marina DEU 46% similarity	3 Phycologia USA 41% similarity	4 Phycological Research USA 39% similarity
---	--	---	--


SJR



Total Cites Self-Cites



External Cites per Doc Cites per Doc




Citable documents Non-citable documents



% International Collaboration



Cited documents Uncited documents

Citations per document





← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scima
```

Metrics based on Scopus® data as of April 2020

M **Muhammad Hendri** 2 years ago

I intend to send my writing / paper entitled:

Identification and Test of Pathogenicity of Bacteria Causes of Ice-ice Disease on *Eucheuma spinosum* Seaweed collected from floating cages from Lampung Bay waters.

This is the abstract.

Seaweed is one of the marine commodities that has many enthusiasts one of which is a type *E. spinosum*. This type of seaweed is in great demand because of its ability to produce carrageenan. However the cultivation of seaweed from the *Eucheuma* genus has encountered many obstacles, both due to physical, chemical and biological factors. The biggest obstacle to seaweed cultivation is disease. One disease that commonly attacks seaweed is ice-ice. This research to determine the number of isolates obtained while identifying the isolate bacteria from *E. spinosum* seaweed which is attacked by ice-ice, with the addition of pathogenicity test to see the level of pathogenicity of each identified bacterium. The research was conducted in October 2018-January 2019 at the Fish Quarantine Center, Palembang and the Center FOR THE Development of Marine Cultivation (BBPBL), Lampung. Bacterial identification is done by method Microbact™ Identification Kit, while the pathogenicity test was carried out using the design of the RAL (complete and random design) for 7 days. The results of the study showed that 5 isolates were identified with bacteria *V. alginolyticus*, *A. hydrophila*, *P. aeruginosa* and *P. myxofaciens*. Pathogenicity test showed that *A. hydrophila* bacteria had the fastest rate of infection which was offset by the highest rate of weight loss during the testing process.

Keywords : *E. spinosum*, bacteria that cause ice-ice disease, pathogenicity test

Regards

Muhammad Hendri

reply




60 articles in this issue

1. Microalgal cultivation in porous substrate bioreactor for extracellular polysaccharide production

Authors

- o Alice Ekelhof
- o Michael Melkonian
- o Content type: OriginalPaper
- o Published: 27 December 2016
- o Pages: 1115 - 1122



2. [Growth and photosynthetic activity of *Botryococcus braunii* biofilms](#)

Authors (first, second and last of 5)


- o Risa S. Wijihastuti
- o Navid R. Moheimani
- o Makoto M. Watanabe
- o Content type: OriginalPaper
- o Published: 20 December 2016
- o Pages: 1123 - 1134



3. [Application of sodium erythorbate to promote the growth of *Chlorella vulgaris*](#)

Authors (first, second and last of 4)


- o Hongwu Cui
- o Fanping Meng
- o Yuejie Wang
- o Content type: OriginalPaper
- o Published: 30 November 2016
- o Pages: 1135 - 1144



4. [Effects of cell motility and morphology on the rheology of algae suspensions](#)

Authors (first, second and last of 5)


- o N. Cagney
- o T. Zhang
- o S. Balabani
- o Content type: OriginalPaper
- o Published: 18 January 2017
- o Pages: 1145 - 1157



5. [Differential growth and biochemical composition of photoautotrophic and heterotrophic *Isochrysis maritima*: evaluation for use as aquaculture feed](#)

Authors


- o E. Mohammad Basri
- o W. O. Wan Maznah
- o Content type: OriginalPaper
- o Published: 19 January 2017
- o Pages: 1159 - 1170



6. Cultivation of *Chlorella vulgaris* with swine wastewater and potential for algal biodiesel production

Authors (first, second and last of 5)


- Kibok Nam
- Hansol Lee
- Jong-In Han
- Content type: OriginalPaper
- Published: 02 December 2016
- Pages: 1171 - 1178



7. Comparison of the culture and harvesting of *Chlorella vulgaris* and *Tetrademus obliquus* for the removal of pharmaceuticals from water

Authors (first, second and last of 5)


- C. Escapa
- R. N. Coimbra
- M. Otero
- Content type: OriginalPaper
- Published: 02 December 2016
- Pages: 1179 - 1193



8. Visualization and quantification of oil in single microalgal cells

Authors (first, second and last of 4)


- Mahmoud Al Ahmad
- Sulaiman Al-Zuhair
- Farah Mustafa
- Content type: OriginalPaper
- Published: 18 November 2016
- Pages: 1195 - 1202



9. Effect of light quality on the growth and proximal composition of *Amphora* sp.

Authors

- Celia Carolina Romero-Romero
- M. del Pilar Sánchez-Saavedra
- Content type: OriginalPaper
- Published: 17 December 2016
- Pages: 1203 - 1211



10. [Progress and challenges in producing polyhydroxyalkanoate biopolymers from cyanobacteria](#)

Authors (first, second and last of 4)

- Akhilesh Kumar Singh
- Laxuman Sharma
- Jyoti Mala
- Content type: ReviewPaper
- Published: 26 November 2016
- Pages: 1213 - 1232

n = 1 R = hydrogen Poly(3-hydroxypropionate)


R = methyl Poly(3-hydroxybutyrate)

R = ethyl Poly(3-hydroxyvalerate)

11. [Thermostable phycocyanin from the red microalga *Cyanidioschyzon merolae*, a new natural blue food colorant](#)

Authors (first, second and last of 5)

- D. Y. Rahman
- F. D. Sarian
- M. J. E. C. van der Maarel
- Content type: OriginalPaper
- Open Access
- Published: 21 November 2016
- Pages: 1233 - 1239



12. [Analysis of phospholipids and neutral lipids in three common northern cold water diatoms: *Coscinodiscus concinnus*, *Porosira glacialis*, and *Chaetoceros socialis*, by ultra-high performance liquid chromatography-mass spectrometry](#)


Authors (first, second and last of 5)

- E.Y. Artamonova
- J.B. Svenning
- H.C. Eilertsen
- Content type: OriginalPaper
- Published: 19 January 2017
- Pages: 1241 - 1249

13. [Analysis of metabolic responses of *Dunaliella salina* to phosphorus deprivation](#)

Authors (first, second and last of 4)


- Hexin Lv
- Xianggan Cui
- Shiru Jia
- Content type: OriginalPaper
- Published: 24 January 2017
- Pages: 1251 - 1260



14. [Ammonium chloride: a novel effective and inexpensive salt solution for phycocyanin extraction from *Arthrospira \(Spirulina\) platensis*](#)

Authors (first, second and last of 7)


- Emmanuel Manirafasha
- Theophile Murwanashyaka
- Keju Jing
- Content type: OriginalPaper
- Published: 09 January 2017
- Pages: 1261 - 1270



15. [A comparison of protein extraction methods optimizing high protein yields from marine algae and cyanobacteria](#)

Authors (first, second and last of 4)


- Lacey M. Field
- Wayne R. Fagerberg
- S. Anne Böttger
- Content type: OriginalPaper
- Published: 06 December 2016
- Pages: 1271 - 1278



16. [Detection of microcystin producing cyanobacteria in Spirulina dietary supplements using multiplex HRM quantitative PCR](#)

Authors (first, second and last of 4)


- Kamath Mukund Manali
- Rex Arunraj
- Mohandass Ramya
- Content type: OriginalPaper
- Published: 16 November 2016
- Pages: 1279 - 1286



17. [Impacts of enhanced UVB radiation on photosynthetic characteristics of the marine diatom *Phaeodactylum tricornutum* \(Bacillariophyceae, Heterokontophyta\)](#)

Authors (first, second and last of 5)


- Kunpeng Shi
- Jia Yu
- Xuexi Tang
- Content type: OriginalPaper
- Published: 01 February 2017
- Pages: 1287 - 1296



18. [Biocidal effect of \(*E*\)-anethole on the cyanobacterium *Aphanizomenon gracile* Lemmermann](#)

Authors (first, second and last of 7)


- Nikoletta Ntalli
- Antonis Michaelakis
- Slawek Cerbin
- Content type: OriginalPaper
- Published: 09 December 2016
- Pages: 1297 - 1305



19. [Cell disruption of *Chlorella vulgaris* using active extracellular substances from *Bacillus thuringiensis* ITRI-G1 is a programmed cell death event](#)

Authors (first, second and last of 6)


- Ming-Der Bai
- Hui-Ju Hsu
- Jen-Chih Chen
- Content type: OriginalPaper
- Published: 24 January 2017
- Pages: 1307 - 1315



20. [Antioxidant responses of different microalgal species to nonylphenol-induced oxidative stress](#)

Authors

- Q. T. Gao
- Y. S. Wong
- Nora F. Y. Tam
- Content type: OriginalPaper
- Published: 01 February 2017
- Pages: 1317 - 1329




21. [Bioremediation efficacy—comparison of nutrient removal from an anaerobic digest waste-based medium by an algal consortium before and after cryopreservation](#)

Authors (first, second and last of 5)

- Alla Silkina
- Graham D. Nelson
- John G. Day
- Content type: OriginalPaper


- Open Access
- Published: 05 February 2017
- Pages: 1331 - 1341



22. [Comparison of bacterioplankton communities between before and after inoculation with an algicidal material, Ca-aminoclay, to mitigate *Cochlodinium polykrikoides* blooms: assessment using microcosm experiments](#)

Authors (first, second and last of 4)


- Seung Won Jung
- Seong Yu Noh
- Taek-Kyun Lee
- Content type: OriginalPaper
- Published: 31 January 2017
- Pages: 1343 - 1354



23. [Determination of the toxicity of the freshwater cyanobacterium *Woronichinia naegeliiana* \(Unger\) Elenkin](#)

Authors


- Beata Bober
- Jan Bialczyk
- Content type: OriginalPaper
- Open Access
- Published: 03 February 2017
- Pages: 1355 - 1362



24. [OvMeter: an automated 3D-integrated opto-electronic system for *Ostreopsis cf. ovata* bloom monitoring](#)

Authors (first, second and last of 6)


- Francesca Sbrana
- Ettore Landini
- Massimo Vassalli
- Content type: OriginalPaper
- Published: 03 February 2017
- Pages: 1363 - 1375




25. [Toward mosquito control with a green alga: expression of Cry toxins of *Bacillus thuringiensis* subsp. *israelensis* \(Bti\) in the chloroplast of *Chlamydomonas*](#)

Authors (first, second and last of 4)


- Seongjoon Kang
- Obed W. Odom
- David L. Herrin
- Content type: OriginalPaper
- Published: 23 November 2016
- Pages: 1377 - 1389

**26. [Equilibrium and kinetic studies of Cu\(II\) and Ni\(II\) sorption on living *Euglena gracilis*](#)****Authors**


- Cameron Winters
- Céline Guéguen
- Adam Noble
- Content type: OriginalPaper
- Published: 24 December 2016
- Pages: 1391 - 1398

**27. [New approach for enhancing *Chlorella vulgaris* biomass recovery using ZnAl-layered double hydroxide nanosheets](#)****Authors (first, second and last of 5)**

- Nada Elgiddawy
- Tamer M. Essam
- Ahmed A. Farghali
- Content type: OriginalPaper
- Published: 11 January 2017
- Pages: 1399 - 1407

**28. [Ongrowing and enhancement of n-3 HUFA profile in adult *Artemia*: short- vs long-time enrichment](#)****Authors (first, second and last of 5)**


- Miquel Planas
- Catarina Silva
- Sonia Piñero
- Content type: OriginalPaper
- Published: 06 December 2016
- Pages: 1409 - 1420



29. [Application of sodium alginate in induced biological soil crusts: enhancing the sand stabilization in the early stage](#)

Authors (first, second and last of 7)

- Chengrong Peng
- Jiaoli Zheng
- Yongding Liu
- Content type: OriginalPaper
- Published: 25 January 2017
- Pages: 1421 - 1428



30. [Effect of water velocity on *Undaria pinnatifida* and *Saccharina japonica* growth in a novel tank system designed for macroalgae cultivation](#)

Authors (first, second and last of 6)


- Yoichi Sato
- Masaki Yamaguchi
- Shigeyuki Kawano
- Content type: OriginalPaper
- Published: 18 November 2016
- Pages: 1429 - 1436



31. [Within-species and temperature-related variation in the growth and natural products of the red alga *Asparagopsis taxiformis*](#)

Authors (first, second and last of 6)


- Leonardo Mata
- Rebecca J. Lawton
- Nicholas A. Paul
- Content type: OriginalPaper
- Published: 06 December 2016
- Pages: 1437 - 1447



32. [Limited evolutionary responses to harvesting regime in the intensive production of algae](#)

Authors (first, second and last of 4)


- Rebecca J Lawton
- Nicholas A Paul
- Keyne Monro
- Content type: OriginalPaper
- Published: 07 January 2017
- Pages: 1449 - 1459



33. [Contrasting effects of blue and red LED irradiances on the growth of *Sargassum horneri* during the germling and immature stages](#)

Authors (first, second and last of 7)


- Osamu Miki
- Chikako Okumura
- Toshiaki Kato
- Content type: OriginalPaper
- Published: 02 December 2016
- Pages: 1461 - 1469



34. [Erratum to: Contrasting effects of blue and red LED irradiances on the growth of *Sargassum horneri* during the germling and immature stages](#)

Authors (first, second and last of 7)

- Osamu Miki
- Chikako Okumura
- Toshiaki Kato
- Content type: Erratum
- Published: 04 February 2017
- Pages: 1471 - 1471



35. [Morphological and genetic differentiation of cultivated *Undaria pinnatifida* \(Laminariales, Phaeophyta\)](#)

Authors (first, second and last of 4)

- Kyosuke Niwa
- Atsushi Kobiyama
- Takashi Sakamoto
- Content type: OriginalPaper
- Published: 06 January 2017
- Pages: 1473 - 1482



36. [Erratum to: Morphological and genetic differentiation of cultivated *Undaria pinnatifida* \(Laminariales, Phaeophyta\)](#)

Authors (first, second and last of 4)


- Kyosuke Niwa
- Atsushi Kobiyama

- Takashi Sakamoto
- Content type: Erratum
- Published: 08 March 2017
- Pages: 1483 - 1483

37. [*Hydrurus foetidus* \(Chrysophyceae\)—an inland macroalga with potential](#)

Authors


- Dag Klaveness
- Content type: OriginalPaper
- Published: 09 January 2017
- Pages: 1485 - 1491



38. [Compositional variations of brown seaweeds *Laminaria digitata* and *Saccharina latissima* in Danish waters](#)

Authors (first, second and last of 5)


- Dirk Manns
- Mette Møller Nielsen
- Anne S. Meyer
- Content type: OriginalPaper
- Published: 28 January 2017
- Pages: 1493 - 1506



39. [Novel rapid method for the characterisation of polymeric sugars from macroalgae](#)

Authors (first, second and last of 5)


- S. E. Spicer
- J. M. M. Adams
- Ana L. Winters
- Content type: OriginalPaper
- Open Access
- Published: 16 November 2016
- Pages: 1507 - 1513



40. [Sequential extraction and characterization of fucoidans and alginates from *Ecklonia radiata*, *Macrocystis pyrifera*, *Durvillaea potatorum*, and *Seirococcus axillaris*](#)

Authors (first, second and last of 7)

- Andrew J. Lorbeer
- Suvimol Charoensiddhi
- Wei Zhang
- Content type: OriginalPaper
- Published: 10 November 2016
- Pages: 1515 - 1526



41. [Factors affecting yield and gelling properties of agar](#)

Authors (first, second and last of 6)


- Wei-Kang Lee
- Yi-Yi Lim
- Chai-Ling Ho
- Content type: ReviewPaper
- Published: 28 November 2016
- Pages: 1527 - 1540



42. [A nanocomposite film fabricated with simultaneously extracted protein-polysaccharide from a marine alga and TiO₂nanoparticles](#)

Authors (first, second and last of 4)


- Qingyan He
- Yan Huang
- Shaoyun Wang
- Content type: OriginalPaper
- Published: 27 December 2016
- Pages: 1541 - 1552



43. [Ultrasound-assisted extraction of fucoidan from *Sargassum muticum*](#)

Authors (first, second and last of 5)

- Noelia Flórez-Fernández
- Marta López-García
- Herminia Domínguez
- Content type: OriginalPaper
- Published: 06 January 2017
- Pages: 1553 - 1561




44. [Mastocarpus stellatus as a source of R-phycoerythrin: optimization of enzyme assisted extraction using response surface methodology](#)

Authors (first, second and last of 4)

- Huu Phuoc Trang Nguyen
- Michèle Moranc ais
- Justine Dumay
- Content type: OriginalPaper

- Published: 03 December 2016


- Pages: 1563 - 1570



45. [Chemical composition of volatile compounds in two red seaweeds, *Pterocladia capillacea* and *Osmundaria obtusiloba*, using static headspace gas chromatography mass spectrometry](#)

Authors (first, second and last of 8)


- Daniel Barroso de Alencar
- Jaécio Carlos Diniz
- Silvana Saker-Sampaio
- Content type: OriginalPaper
- Published: 26 November 2016
- Pages: 1571 - 1576



46. [The effects of processing on the in vitro antimethanogenic capacity and concentration of secondary metabolites of *Asparagopsis taxiformis*](#)

Authors (first, second and last of 5)


- Matthew J. Vucko
- Marie Magnusson
- Rocky de Nys
- Content type: OriginalPaper
- Published: 15 November 2016
- Pages: 1577 - 1586



47. [Identification and large isolation of an anti-inflammatory compound from an edible brown seaweed, *Undariopsis peterseniana*, and evaluation on its anti-inflammatory effect in in vitro and in vivo zebrafish](#)

Authors (first, second and last of 11)


- Ji-Hyeok Lee
- Ju-Young Ko
- You-Jin Jeon
- Content type: OriginalPaper
- Published: 24 November 2016
- Pages: 1587 - 1596



48. [Protein extract from red seaweed *Gracilaria fisheri* prevents acute hepatopancreatic necrosis disease \(AHPND\) infection in shrimp](#)

Authors (first, second and last of 4)


- Nantavadee Boonsri
- Tawut Rudtanatip
- Kanokpan Wongprasert
- Content type: OriginalPaper
- Published: 17 November 2016
- Pages: 1597 - 1608



49. [Characteristics of functional materials recovered from Solomon Islands red seaweed \(*Kappaphycus alvarezii*\) using pressurized hot water extraction](#)

Authors (first, second and last of 4)


- Collin Rudolf Nobbs Gereniu
- Periaswamy Sivagnanam Saravana
- Byung-Soo Chun
- Content type: OriginalPaper
- Published: 23 January 2017
- Pages: 1609 - 1621



50. [Anti-proliferation peptides from protein hydrolysates of *Pyropia haitanensis*](#)

Authors (first, second and last of 4)

- Xinliang Mao
- Lu Bai
- Xuewu Zhang
- Content type: OriginalPaper
- Published: 04 January 2017
- Pages: 1623 - 1633



Previous page

-
- 1
- [2](#)

[Skip to main content](#)

Advertisement



[Search](#) 

- [Authors & Editors](#)
- [My account](#)

Menu

- [Authors & Editors](#)
- [My account](#)



[Journal of Applied Phycology](#)

- [Journal home](#) >
- [Editors](#)

Editors

EDITOR-IN-CHIEF

Michael A. Borowitzka

School of Biological Sciences & Biotechnology, Murdoch University, WA, Australia

FOUNDING EDITOR

Brian A. Whitton

University of Durham, UK

ASSISTANT EDITORS

Alejandro H. Buschmann, Universidad de Los Lagos, Centro i~mar and CeBiB, Puerto Montt, Chile

EonSeon Jin, Hanyang University, Seoul, Korea

Stefan Kraan, The Seaweed Company, Mausrevagh, kilcoona, Headford, Ireland

Phang Siew Moi, Institute Of Ocean And Earth Sciences, University of Malaya, Malaysia

Roberto de Philippis, University of Florence, Italy

Alexei Solovchenko, Lomonosov Moscow State University, Russia

Avigad Vonshak, Ben-Gurion University of the Negev, Beer-Sheva, Israel

EDITORIAL BOARD

Rhodora V. Azanza, University of the Philippines, Diliman, Philippines

Sung Min Boo, Chungnam National University, Daejeon, Korea
Steven Feng Chen, College of Engineering, Peking University, Beijing China
Yusuf Chisti, Massey University, Palmerston North, New Zealand
Alan Critchley, Cape Breton University, Nova Scotia, Canada
Joël Fleurence, University of Nantes, Nantes, France
Kunshan Gao, Xiamen University, Xiamen, Fujian, China
Freddy Guilhéneuf, Observatoire Océanologique de Villefranche sur Mer, France
Gustaaf Hallegraeff, University of Tasmania, Hobart, Australia
Ignacio Hernandez Carrero, Universidad de Cadiz, Puerto Real, Cadiz, Spain
Quiang Hu, Institute of Hydrobiology, Wuhan, China
Inna Khozin-Goldberg, Ben-Gurion University of the Negev, Beer-Sheva, Israel
Marc Lahaye, INRA, Nantes, France
Choul-Gyun Lee, Inha University, Department of Biological Engineering, Icheon, Korea
Ira Levine, University of Southern Maine, Portland, ME, USA
Brett Neilan, The University of Newcastle, NSW, Australia
Amir Neori, Israel Oceanographic & Limnological Research, National Center for Mariculture, Eilat, Israel
Shao Jun Pang, Institute of Oceanology, CAS, Qindao, China
Nicholas Paul, University of the Sunshine Coast, Sippy Downs, QLD, Australia
Jürgen E.W. Polle, Brooklyn College, The City University of New York, USA
Ondřej Prášil, Czech Academy of Sciences, Centre ALGHATECH, Trebon, Czech Republic
C.R.K. Reddy, Institute of Chemical Technology, Mumbai India
Rossita Shapawi, Borneo Marine Research Institute, Universiti Malaysia Sabah, Malaysia
Dagmar Stengel, National University of Ireland Galway, Ireland
Wendy A. Stirk, University of KwaZulu-Natal Pietermaritzburg, South Africa
Judy Sutherland, National Institute for Water and Atmospheric Research (NIWA), New Zealand
Milan Szabo, Hungarian Academy of Sciences, Szeged, Hungary
Mario R. Tredici, University of Florence, Italy
Luisa Valente, CIIMAR, Matosinhos, Portugal
Dries Vandamme, University of Hasselt, Belgium
Makoto Watanabe, University of Tsukuba, Ibaraki, Japan
R. Wijffels, Wageningen University, Wageningen, The Netherlands
Nair Yokoya, Instituto de Botanica, Sao Paulo, SP, Brazil
Wei Zhang, Flinders University, Bedford Park, SA, Australia

For authors

[Submission guidelines](#) [Ethics & disclosures](#) [Fees and funding](#) [Contact the journal](#) [Submit manuscript](#)

Explore

[Online first articles](#) [Volumes and issues](#) [Sign up for alerts](#)

Advertisement



Publish with us

[Authors & Editors](#)

[Journal authors](#)

About this journal

Electronic ISSN
1573-5176

Print ISSN
0921-8971

Abstracted and indexed in

1. AGRICOLA
2. BFI List
3. BIOSIS
4. Biological Abstracts
5. CAB Abstracts
6. CNKI
7. Chemical Abstracts Service (CAS)
8. Current Contents/ Agriculture, Biology & Environmental Sciences
9. Dimensions
10. EBSCO Academic Search
11. EBSCO Discovery Service
12. EMBiology
13. Google Scholar
14. Institute of Scientific and Technical Information of China
15. Japanese Science and Technology Agency (JST)
16. Journal Citation Reports/Science Edition
17. Meta
18. Naver
19. OCLC WorldCat Discovery Service
20. ProQuest Agricultural & Environmental Science Database
21. ProQuest Algology, Mycology and Protozoology Abstracts (Microbiology C)
22. ProQuest Aquatic Sciences and Fisheries Abstracts (ASFA)
23. ProQuest Biological Science Database
24. ProQuest Central
25. ProQuest Earth, Atmospheric & Aquatic Science Database
26. ProQuest Natural Science Collection
27. ProQuest Oceanic Abstracts
28. ProQuest SciTech Premium Collection
29. ProQuest-ExLibris Primo
30. ProQuest-ExLibris Summon
31. SCImago
32. SCOPUS
33. Science Citation Index
34. Science Citation Index Expanded (SciSearch)
35. TD Net Discovery Service
36. UGC-CARE List (India)
37. Zoological Record

Copyright information

[Rights and permissions](#)

[Springer policies](#)

Growth and photosynthetic activity of *Botryococcus braunii* biofilms

Risa S. Wijihastuti¹ · Navid R. Moheimani¹ · Parisa A. Bahri² ·
Jeffrey J. Cosgrove¹ · Makoto M. Watanabe³

Received: 30 August 2016 / Revised and accepted: 12 December 2016
© Springer Science+Business Media Dordrecht 2016

Abstract *Botryococcus braunii* is a green microalga capable of producing large amounts of external long-chain hydrocarbons suitable as a source of biofuel. There have been several studies indicating that cultures of *B. braunii* can reduce the energy and water requirement for mass biofuel production, especially if non-destructive extraction methods for milking hydrocarbons are used. Growing microalgae as a raw material for biofuel using conventional liquid-based cultivation (i.e., raceway ponds) has yet to be shown to be economically successful. An alternative solid growth (biofilm) cultivation method can markedly reduce the energy requirements and costs associated with the harvesting and dewatering processes. We evaluated the growth of biofilms of several strains of *B. braunii* (from races A, B, L and S) and found that three of the four tested races successfully grew to stationary phase in 10 weeks with no contamination. Among all races, *B. braunii* BOT22 (race B) reached the highest biomass and lipid yields (3.80 mg dry weight cm⁻² day⁻¹ and 1.11 mg dry weight cm⁻²). Irrespective of the race, almost all photosynthetic parameters (F_v/F_0 , PI_{ABS} and the OJIP curve) showed that the biofilm cultures were more stressed during lag and stationary phases than in logarithmic phase. We also studied the *Botryococcus* biofilm profiles using confocal microscopy

and found that this method is suitable for estimating the overall biomass yield when compared with gravimetric measurement. In conclusion, the growth characteristics (biomass and lipid) and photosynthetic performance of all races indicated that *B. braunii* BOT22 is the most promising strain for biofilm cultivation.

Keywords Microalgae · Chlorophyta · Biofuel · Productivity · Hydrocarbon · Quantum yield · OJIP

Introduction

Biofuel has been proposed for many years as a renewable source of energy to reduce the existing demand for fossil fuels (Borowitzka 2013; Borowitzka and Moheimani 2013). Microalgae are claimed to be excellent candidates for biofuel production (Dixon 2013), as they can have high lipid contents (Fon Sing et al. 2013) and can be grown on non-arable land using wastewater (Graham et al. 2009; Dixon 2013). However, the costs of microalgal cultivation and downstream processing to yield the final biofuel product remain high (de Boer et al. 2012). Some of the main reasons for the high cost of production include harvesting-dewatering and extraction of lipids (Scott et al. 2010; Fon Sing et al. 2013). While innovation in downstream processing is required to make the production of microalgae biofuel economically feasible, improvements in cultivation are also necessary. Liquid-based cultivation has been the most common method for the mass culture of microalgae (Borowitzka 2013). The vast amounts of water and nutrients demanded by liquid-based cultivation processes are an issue for microalgal-based commodity products such as biofuels (Borowitzka and Moheimani 2013). An alternative way to cultivate microalgae is to grow them as a biofilm (Berner et al. 2015). Liquid cultures generally have a biomass

✉ Navid R. Moheimani
n.moheimani@murdoch.edu.au

¹ Algae R&D Centre, School of Veterinary and Life Sciences, Murdoch University, Murdoch, WA 6150, Australia

² School of Engineering and Information Technology, Murdoch University, Murdoch, WA 6150, Australia

³ Algae Biomass and Energy System R&D Center (ABES), University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan

content of between 0.02 and 0.06% solids (Schnurr et al. 2013). However, in a biofilm the organic solids can be as high as 16% (Ozkan et al. 2012). Cultivation of microalgae for biofuel in liquid has three main stages: growth of biomass, biomass harvesting-dewatering and lipid extraction. Biofilm growth can significantly reduce water consumption and dewatering requirements, as well as minimising nutrient use while improving light capture (Liu et al. 2013; Berner et al. 2015). Ozkan et al. (2012) reported that that biofilm cultivation can reduce water usage per kilogramme of algal biomass by 45% and the dewatering energy requirement by 99.7% compared to open pond cultivation. Therefore, the development of large-scale biofilm growth systems for microalgae such as *Botryococcus braunii* can potentially overcome some of the challenges of microalgal biofuel production.

The green microalga *B. braunii* produces significant amounts of long-chain unbranched hydrocarbons (Banerjee et al. 2002; Metzger and Largeau 2005). This species is divided into four races based on the type of hydrocarbon produced (Kawachi et al. 2012). In some cases, especially when cells are in very late stationary phase, the hydrocarbon content of *B. braunii* can be as high as 86% of the organic weight (Brown et al. 1969; Dayananda et al. 2007). These hydrocarbons can be converted to transport fuels due to their similarity to fossil fuel (Hillen et al. 1982). Importantly, *B. braunii* accumulates these hydrocarbons in the extracellular matrix, (Banerjee et al. 2002; Eroglu and Melis 2010) and destructive extraction to harvest the hydrocarbons can potentially be avoided when using this species to produce biofuel if non-destructive hydrocarbon extraction techniques are applied (Moheimani et al. 2013, 2014). By maintaining viable biomass after harvesting, non-destructive hydrocarbon extraction, also known as algal milking, substantially reduces the importance of algal growth rate and increases hydrocarbon productivity (Moheimani et al. 2013, 2014). Chaudry et al. (2015) have shown that non-destructive hydrocarbon production from *B. braunii* consumes 70% less energy and 30% less water compared to conventional wet lipid extraction. This extraction method has been proven as an energetically feasible process for hydrocarbon production from *B. braunii* combined with open pond cultivation, cylindrical sieve rotator filter system dewatering and nanofiltration as the solvent recovery stages (Chaudry et al. 2017). Therefore, non-destructive hydrocarbon extraction from *B. braunii* can be attractive for sustainable biofuel production.

Current knowledge of *B. braunii* biofilm growth and lipid productivity is limited. Therefore, in the present work, we assessed the potential of biofilm growth and the overall lipid productivity of different *B. braunii* races (A, B, L and S). Considering that microalgal biomass productivity is closely

correlated with photosynthesis, photosynthetic performance of all strains was also measured.

Materials and methods

Microalgae culture source and liquid culture maintenance

Four strains of *B. braunii*, one from each race, were used in this study. *Botryococcus braunii* strains BOT22 (race B), BOT84 (race L) and BOT7 (race S) were obtained from the Algae Biomass and Energy System R&D Center (ABES), University of Tsukuba, Tsukuba, Japan. *Botryococcus braunii* UTEX2441 (race A) was obtained from the University of Texas culture collection. All cultures were maintained in modified AF-6 medium (Watanabe et al. 2000) and were grown at 25 ± 2 °C under 50 ± 5 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ provided by natural cool white fluorescent lights with a 12:12 day/night cycle.

Biofilm cultivation

To grow *B. braunii* strains in biofilm, choosing the right nutrient delivery method is essential. Preliminary studies showed that use of a sponge for maintaining constant wetness and medium supply is more effective than agar or adding a liquid medium directly to the biofilm (unpublished data, see online resource 1). In this study, filtering the liquid *B. braunii* culture on to an inoculum attachment surface (2.834 cm^2) was chosen as the inoculation technique (Liu et al. 2013). Various types of material were used to identify the best attachment surface for the *B. braunii* biofilm. Of the five tested materials [glass microfibre (GF/C), cellulose nitrate (CN) and cellulose ester filter paper; toilet hand towels and Harris™ coffee filter paper], *B. braunii* grew best on cellulose nitrate filter papers (see online resource 2).

Botryococcus braunii were grown in biofilm using CN membrane filters with a $0.45 \mu\text{m}$ pore size and a diameter of 25 mm. The initial liquid stock culture for all strains for inoculation was grown in the same condition for 4 weeks. Five millilitre of the *B. braunii* liquid culture were filtered through the CN filters and these were then placed inside six-well cell culture plates which covered with the lid on top of $35 \text{ mm} \times 10 \text{ mm}$ (diameter \times depth) polyvinyl alcohol (PVA) sponges (media preserver). To start, 8 mL of AF-6 medium was added into each well containing a PVA sponge. One CN membrane filter was placed on the top of each wet sponge. Membrane filters were incubated overnight before being weighed as a blank filter. Pre-weighed membrane filters were inoculated by filtering 5 mL of *B. braunii* stock culture for all strains. We chose 5 mL as the preliminary results indicated this to be the best amount for inoculation of CN filters

(unpublished data). The membrane filters were incubated overnight before being weighed (day 1 of the trials). The biofilm cultures were grown for 10 weeks under an irradiance of $75 \pm 2 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ using natural cool white fluorescent lights with a 12:12 day/night cycle, at $25 \pm 2^\circ\text{C}$. The medium was changed every week only for the first 4 weeks. This provided the biofilm cultures with the same nutrients as 32 mL of the liquid AF-6 medium. To counter evaporative loss, after 4 weeks, sterile deionised water was added to the sponges. Choosing samples for biochemical, photosynthetic activity and biofilm structure analysis was by using a table of random numbers. Filters were photographed weekly using a digital camera for monitoring visual changes in the biofilm cultures.

Analytical measurements

Growth measurement A method for measuring biofilm growth non-destructively was developed by adapting several different weight conversion methods previously used by Gates et al. (1982) and Ricciardi and Bourget (1998). Five different volumes (1, 3, 5, 7 and 10 mL) of *B. braunii* cultures were filtered onto different pre-weighed GF/C filters (each volume had six replicates). The wet weights of the filter papers with algae were measured three times, every 2 h and once after 24 h. The dry weights were determined following Moheimani et al. (2013) after samples were dried overnight at 60°C . Linear regression with the model forced through the origin was performed using Microsoft Excel to determine the correlation between dry weight and wet weight. This analysis resulted in equations for converting wet weight to dry weight, which were applied to allow non-destructive measurements of growth for each race of *B. braunii*. Wet weights were measured weekly using a calibrated five-digit Mettler-Toledo AB135-S balance.

Biofilm structure analysis A fresh sample of algal biofilm was used for confocal microscopy. The biofilm was stained with BODIPY 505/515 (4,4-difluoro-1,3,5,7-tetramethyl-4-bora-3a,4a-diaza-s-indacene, Life Technologies Molecular Probes) to highlight lipid content (Brennan et al. 2012; Cirulis et al. 2012; Govender et al. 2012). *Botryococcus braunii* cells were identified via autofluorescence of chlorophyll content. The BODIPY stain was freshly prepared just prior to observation (Govender et al. 2012) by adding $100 \mu\text{g mL}^{-1}$ BODIPY 505/515 in 2% dimethyl sulfoxide (DMSO), then diluting to $0.75 \mu\text{g mL}^{-1}$ in 2% DMSO, and storing in a dark bottle to avoid light exposure. One drop of the stain was placed on the sample, followed by 4 min incubation in the dark before viewing.

The three-dimensional structure of the biofilms was viewed with a Nikon C2+ multispectral laser scanning confocal microscope using 20 \times objectives (Lawrence et al. 1998; Neu

et al. 2004) and laser excitation at 640.0 nm for observing chlorophyll content and 488.0 nm for observing the BODIPY-stained lipid. Images were taken and processed by the Nikon Imaging Software (NIS) Elements Advanced Research package and processed into three-dimensional composite images approximately $632 \mu\text{m (L)} \times 632 \mu\text{m (W)}$ with various biofilm depths (36–192 μm). Quantitative analyses of the distribution of algal cells and lipid within the biofilm matrix were carried out using COMSTAT software (Heydorn et al. 2000).

Photosynthesis measurement Photosynthesis measurements were performed directly on *B. braunii* biofilms using a Handy-PEA chlorophyll fluorometer (Hansatech Instruments, UK) paired with PEA Plus V1.10 software. This instrument has a high-intensity LED array (3 lamps, centred on 650 nm) with an NIR short-pass filter and allows measurement of the so called *fast phase* of the fluorescence induction curve and estimation of the OJIP parameters. Saturation pulse measurements performed on dark-adapted samples allowed derivation of the photosynthetic parameters. The maximum ratio of quantum yields of photochemical and concurrent non-photochemical processes in PS-II related to the dark adaptation state (F_v/F_0) was one of the parameters that was measured, instead of measuring the more usual maximum quantum yield of PS-II photochemistry (F_v/F_m). This parameter was used because it is a more sensitive parameter than F_v/F_m for indicating the maximum efficiency of photochemical processes in PS-II (Roháček 2002). The other parameters measured were the photosynthesis performance index (PI_{ABS}), the light absorption flux for PS-II antenna chlorophyll (ABS/RC), and energy dissipation at the level of antenna chlorophyll at Time 0 in PS-II (DI_0/RC), all which are the key parameters for the OJIP parameters according to Dao and Beardall (2016). These derivations were performed as described in Cosgrove and Borowitzka (2010) and Strasser et al. (2000) (see the details in online resource 3). Due to the low growth rate of *B. braunii* in the biofilms, the aforementioned photosynthetic measurements (F_v/F_0 , PI_{ABS} , ABS/RC , and DI_0/RC) were measured every 2 weeks. However, the fast phase-fluorescence induction curves were only analysed for three different weeks to represent the chlorophyll fluorescence transient changes in each growth phase, which were the lag phase (week 0), the logarithmic phase (week 6), and the stationary phase (week 10). Along with the fast phase fluorescence induction curve analysis, the relative variable fluorescence at 2 ms (V_J) was also measured to express the connection between PS-II units (Force et al. 2003).

Biofilms were dark-adapted for 20 min prior to measurement to allow re-oxidation of all PS-II reaction centres and estimation of the minimum fluorescence yield. Measurements ($n = 4$) were performed by exposing the samples to a 1.2 s

pulse of high intensity light ($3500 \mu\text{mol photon m}^{-2} \text{s}^{-1}$) to saturate the photosystems and create maximal fluorescence yield. Handy-PEA provided fluorescence values every $10 \mu\text{s}$ to 1ms between 0 and 1.2 s measurement times in logarithmic time sequence. The emitted fluorescence from the sample was recorded and digitised by the Handy-PEA. The data were analysed and displayed graphically using SigmaPlot Version 13 software.

Statistical analysis

One way ANOVA and Tukey's HSD post hoc tests were used to determine the significant difference in the photosynthetic activity during the growth period. The statistical analysis was conducted for each *B. braunii* race.

Results


Four strains, one from each race of *B. braunii*, were grown for 10 weeks as a biofilm (Fig. 1). As *B. braunii* UTEX2441 (race A) cultures were heavily contaminated by fungi from week 5, we stopped recording growth measurements for this strain (Figs. 1 and 2). *Botryococcus braunii* UTEX2441 biofilm was re-grown one more time. However, the culture was again contaminated after week 5. The other three *B. braunii* strains grew well as biofilms, with no contamination between weeks 0 (lag phase) and 10 (stationary phase) (Fig. 2). Furthermore, no contamination was observed in the biofilms up to week 27. Before becoming contaminated, *B. braunii* UTEX2441 showed growth between weeks 1 and 5 (Fig. 2). The other

three strains has a lag phase of three to 5 weeks before entering the logarithmic phase, which was characterised by a substantial increase in biofilm wet weight (Fig. 2).

Botryococcus braunii BOT22 (race B), BOT84 (race L) and BOT7 (race S) reached stationary phase in week 8 (Fig. 2), and the culture colour changed from green to yellowish due to accumulation of carotenoids (Fig. 3). Growth was gravimetrically measured as wet weigh, and this was converted to dry weight using the equations shown in (able 1. Biomass and lipid yields, as well as the productivities of each strain, are summarised in Table 2. Based on the wet weight measurement, *B. braunii* BOT 22 achieved the highest biomass yield ($107.24 \pm 7.86 \text{ mg wet weight cm}^{-2}$) and productivity ($40.61 \pm 25.10 \text{ mg wet weight cm}^{-2} \text{day}^{-1}$), compared to *B. braunii* BOT7 and BOT84. *Botryococcus braunii* BOT22 also showed the highest biomass productivity ($3.80 \pm 2.35 \text{ mg dry weight cm}^{-2} \text{day}^{-1}$) in the logarithmic phase compared to the other strains. *Botryococcus braunii* BOT22 achieved the highest lipid yield, which was $1.11 \pm 0.08 \text{ mg cm}^{-2}$ in the stationary phase (Table 1); while both *B. braunii* BOT7 and BOT84 yielded $0.83 \pm 0.08 \text{ mg cm}^{-2}$ and $0.83 \pm 0.03 \text{ mg cm}^{-2}$, respectively. However, among the tested strains the highest lipid content (26.6%) was achieved in the culture of *B. braunii* BOT7 (Table 2).

Biofilm thickness for all strains increased over time, ranging from 62 to 92 μm from the initial thickness between weeks 0 and 10 (Figs. 3 and 4). The area under the curve for the cell and lipid distributions of *B. braunii* are presented in each graph (Fig. 4). Both the cell and the lipid distribution areas increased and reached their peak between 20 and 60 μm from the biofilm surface, and then gradually decreased in the deeper

Fig. 1 *Botryococcus braunii* biofilm cultures during the lag phase (week 0), logarithmic phase (week 6) and stationary phase (week 10). Each culture is 25 mm in diameter






Fig. 2 Growth and photosynthetic activity of all *Botryococcus braunii* biofilm cultures during the growth period. The growth curve based on the wet weight (*error bars* = standard deviation); photosynthesis performance index (PI_{ABS}), maximum photochemical efficiency (F_v/F_0) (dark adapted); the light absorption flux for PSII antenna chlorophyll

(ABS/RC); the dissipation at the level of antenna chlorophyll at time 0 in PS-II (DI_0/RC). The *same letter above each bar*, indicates no significant difference (one way ANOVA, $P > 0.05$, $n = 4$). For the photosynthetic activity measurements, the *error bars* indicate the range

layers of the biofilm (Fig. 4), even though the biofilm thickness increased during the growth period. Based on the area

under the curve measurements, the cell and lipid distribution areas of *B. braunii* BOT7 and BOT22 showed a similar




Fig. 3 Confocal microscope images of *Botryococcus braunii* BOT 7 (a, b, c), BOT 22 (d, e, f), and BOT 84 (g, h, i) in the lag (week 0) (a, d, g), logarithmic (week 6) (b, e, h) and stationary (week 10) (c, f, i) phase

pattern, increasing from between lag phase and log phase and then decreasing by the beginning of the stationary phase (Fig. 4). On the other hand, both the cell and lipid distribution areas of *B. braunii* BOT84, which had a bigger initial cell distribution in the lag phase, increased through the log and stationary phases of growth.

The confocal microscopy analysis of algal cell distribution showed a similar trend to the dry weight measurement results. The highest productivity was achieved in the biofilm of *B. braunii* BOT22. The ratio of total area that covered by the lipid to the cells (the lipid ratio of the cells) decreased during the logarithmic phase and increased in the stationary phase in all three *B. braunii* strains. However, in *B. braunii* BOT22 this ratio increased greatly compared to the other strains from the logarithmic to the stationary phase.

Table 1 The equations for converting wet weight (x) into dry weight (y) for all *B. braunii* strains

Strain	Equation	R ²
UTEX2441	$y = 0.17x$	0.85
BOT7	$y = 0.04x$	0.93
BOT22	$y = 0.09x$	0.93
BOT84	$y = 0.22x$	0.68

Chlorophyll fluorescence parameters were used to rapidly and non-destructively investigate the relationship between biofilm growth and photosynthetic activity in the *B. braunii* strains (Fig. 2). Although the growth curves showed that most *B. braunii* strains remained in lag phase until week 5 the photosynthetic activity parameters (PI_{ABS} , F_v/F_0 , ABS/RC and DI_0/RC) showed increases from week 2. Lower PI_{ABS} and F_v/F_0 were found in week 0 than in week 2, indicating that *B. braunii* strains were stressed at the beginning of the experiment (Fig. 2). The ABS/RC ($F(2.81) = 74.764$ (BOT7); 56.147 (BOT22); 185.363 (BOT84), $P < 0.05$) and DI_0/RC ($F(2.81) = 37.688$ (BOT7); 49.380 (BOT22); 120.786 (BOT84), $P < 0.05$) decreased significantly from week 0 to week 2 as the algae began to acclimate to the new environment (Fig. 2). The PI_{ABS} values for each of the remaining strains remained steady for *B. braunii* BOT7 and increased for *B. braunii* BOT22 and BOT84 when they entered the logarithmic phase between weeks 2 and 6. The same phenomenon was observed for F_v/F_0 , indicating that photosynthetic activity improved when cells began entering logarithmic phase in week 6. The PI_{ABS} and F_v/F_0 values of *B. braunii* UTEX 2441 also increased in week 2 but then decreased at week 4 due to the

Table 2 Biomass yield, productivity, lipid yield of *B. braunii* biofilm

Strain	Race	Yield (mg cm ⁻²)		Productivity* (mg cm ⁻² day ⁻¹)		Lipid yield (mg dry weight cm ⁻²)
		Wet weight	Estimated dry weight**	Wet weight	Estimated dry weight**	
BOT7	S	72.44 ± 7.68	3.12 ± 0.33	9.76 ± 1.63	0.42 ± 0.07	0.83 ± 0.08
BOT22	B	107.24 ± 7.68	10.04 ± 0.74	40.61 ± 25.10	3.80 ± 2.35	1.11 ± 0.08
BOT84	L	62.31 ± 3.62	13.60 ± 0.79	4.53 ± 0.77	0.99 ± 0.17	0.83 ± 0.03

*Productivity was calculated during the logarithmic phase (weeks 5–6 for BOT2, weeks 5–8 for BOT84 and BOT7)

**Based on the equations on Table 1

(data are means ± standard deviation, $n = 6$)

contamination load (Fig. 2). Among the strains that survived the 10 weeks and reached stationary phase (week 8), the PI_{ABS} and F_v/F_0 values were observed to decrease, except for the PI_{ABS} value of *B. braunii* BOT22, which was still increasing. Within each strain, the recorded ABS/RC did not change significantly in the logarithmic phase until the culture reached stationary phase. Similarly, the DI_0/RC values showed no significant change during logarithmic and stationary phases, except for *B. braunii* BOT84.

The fast phase of fluorescence induction curves (OJIP), which were measured in three different growth phases, revealed similar patterns to those of the other photosynthetic activity parameters (Fig. 5). In the lag phase (week 0), the biofilm was apparently under stress, as indicated by the appearance of a large inflexion on the curve (Fig. 5) and also the higher V_J compared to other growth phase (Fig. 6). Photosynthetic activity improved in the logarithmic phase (week 6), which was characterised by reduced inflexion and V_J (Fig. 5). There were no obvious inflexions on the curve for *B. braunii* BOT7, BOT22 and BOT84 in the stationary phase compared to the previous growth phase, while the V_J values were still decreased in this phase.

Discussion

This study clearly showed that it is possible to grow certain strains of *B. braunii* (three races: B, L and S) as a biofilm with no contamination from the lag phase through to the stationary phase. Our study also indicated that the ability of *B. braunii* to grow in a biofilm is strain-specific. Our *B. braunii* race A cultures became contaminated, and their growth was not sustainable. Interestingly, Metzger and Casadevall (1989) showed that *B. braunii* races B and L are more resistant to contamination due to their high terpenoid content acting as an antimicrobial agent. Although in this study, *B. braunii* race A was less resistant

to contamination, several other studies have used this race in their experiments. However, those studies were conducted over shorter periods than this study. Compared to *B. braunii* liquid cultures grown in open raceway ponds, this study potentially achieved higher biomass productivity. The biomass productivity was higher than the 3.42 mg cm⁻² day⁻¹ recorded by Ashokkumar and Rengasamy (2012) and the 0.75 mg cm⁻² day⁻¹ recorded by Zhang (2013). Ozkan et al. (2012) successfully grew *B. braunii* (LB 572; race A) as a biofilm for 5 weeks with 0.071 ± 0.06 mg DW cm⁻² day⁻¹ biomass productivity (2750 cm² *B. braunii* biofilm at 55 μmol photons m⁻² s⁻¹ resulted in 2.49 ± 0.21 mg DW cm⁻² biomass yield and with 27 ± 2% lipid content). Cheng et al. (2013) grew *B. braunii* biofilm in attached cultivation photobioreactors. The attached photobioreactor produced 0.65 mg DW cm⁻² day⁻¹ biomass productivity (= 6.20 mg DW cm⁻² biomass yield with 42.5% lipid content on 800 cm⁻² surface area under 100 μmol photons m⁻² s⁻¹). Our results indicated a five- and 53-fold higher biomass yield and productivity compared to those reported by Ozkan et al. (2012). *Bortyococcus braunii* BOT22 also showed a two- and sixfolds higher biomass yield and productivity compared to Cheng et al. (2013). However, the *B. braunii* lipid yield achieved in current study was lower than in the Cheng et al. (2013) study. For generating the biofilm, we used sponges that were submerged in a liquid medium similar to the two layer system of Shi et al. (2007), while in other similar studies the medium flowed over the biofilm (Berner et al. 2015). Liquid flow can be critical factors in algal biofilm development (Berner et al. 2015). The static growth method used in our study possibly had a negative influence on the overall lipid production of *B. braunii*.

The multi-layer vertical *B. braunii* biofilm culture used by Cheng et al. (2013) yielded 4.91 mg DW cm⁻² day⁻¹ productivity (56 mg DW cm⁻² biomass yield and 51.6% lipid content). These figures are significantly higher than the results obtained in single-layer horizontal biofilms.




Fig. 4 **a** Growth curve of *B. braunii* strains based on confocal microscope observation time; comparison of *B. braunii* BOT7, BOT22 and BOT84 biofilm thickness; cells and lipid distribution; and the average of area under the curve in: **b** the lag (week 0); **c** logarithmic (week 6); **d** and stationary (week 10) phase. The number of repetitions in measurement was based on sample availability, ranging between 1 and 3

Liu et al. (2013) also used a multi-layer vertical biofilm culture with *Scenedesmus obliquus*, resulting in 7.09 mg DW cm⁻² day⁻¹ biomass productivity (80 mg DW cm⁻² biomass yield and 47% lipid content).

Confocal microscopy also showed that the increased biofilm thickness during the growth period did not increase the cell distribution area in the biofilm. Since the confocal microscope captured the presence of algal cells through chlorophyll autofluorescence (Satpati and Pal 2015), the colour change in the stationary phase biofilm cultures may also reflect the cell distributions observed by confocal microscopy. Colour changes in algal colonies are closely related to the physiological state of the organism (Metzger et al. 1985). *Botryococcus braunii* cells dominated the top layer of biofilm (20–60 µm from the surface) which indicated that in the biofilms, *B. braunii* cells grew on top of each other towards the light (phototropic positive). Therefore, the older cells and colonies dominated the bottom layer. Increasing biofilm thickness can affect algal cells due to light or nutrient limitations in the deeper regions (Berner et al. 2015). As a result of this environmental stress (lack of nutrients) in the stationary phase, the *B. braunii* colour changed from green to yellowish due to an accumulation of carotenoids (Metzger et al. 1985).

Confocal imaging and the profile analysis also indicated that most of lipids were concentrated in the top layer (between the 20–60 µm depth range of the biofilm). This could potentially be very advantageous, especially if the aim is to extract the oils non-destructively. This means

that if solvents are to be used for milking, they can reach the hydrocarbons (the product of interest) first, which may result in fewer detrimental effects on the *B. braunii* cultures. Obviously, the proposed milking method will need to be tested on *B. braunii* cultures grown in biofilm.

Measuring photosynthetic activity is one of the methods to monitor microalgal growth (Consalvey et al. 2005). To the best of our knowledge, this is the first detailed study of photosynthetic activity in *B. braunii* biofilms. *Botryococcus braunii* BOT7, BOT22 and BOT84 showed a similar pattern for all photosynthetic parameters measured. At the start of the experiments, the algal biofilms were stressed because the algae had been moved to a new environment with less water, which changed the nutrient availability and had a higher light intensity (Ozkan et al. 2012; Schnurr et al. 2013; Berner et al. 2015). The reduced water availability and higher light intensity in the biofilm cultivation system resulted in low PI_{ABS} and F_v/F_0 . PI_{ABS} values, which represent the vitality of photosynthetic organisms and cover three primary steps in photosynthetic activity: absorbing light energy, trapping this energy and converting the energy via electron transport (Strauss et al. 2006). The photochemical efficiency (F_v/F_0) represents the ability of the algae to use light for photosynthesis and shows the effect of stressful conditions on an organism's photosynthetic apparatus (Cosgrove and Borowitzka 2010; Dao and Beardall 2016). The low PI_{ABS} and F_v/F_0 values indicate that the algae were under stress due to their new environment. The higher irradiance received by the biofilm system compared to liquid cultures also resulted in a high absorbance of photons per reaction centre (ABS/RC), which reflected the apparent antenna size of the photosynthesis organelle (Demetriou et al. 2007). This phenomenon was also observed by Lu and Vonshak (1999) in their study of photoinhibition caused by high sun irradiance in outdoor




Fig. 5 Fluorescence induction curve for *B. braunii* BOT7, BOT22 and BOT84 in the lag (week 0), logarithmic (week 6) and stationary (week 10) phase




Fig. 6 Relative variable fluorescence at 2 ms (V_J) of *B. braunii* BOT7, BOT22 and BOT84 in the lag (week 0), logarithmic (week 6) and stationary (week 10) phase

Spirulina platensis cultures. The high value of ABS/RC could result from inactivation of the reaction centres (Strasser et al. 1999). At this stage, the low F_V/F_0 and the high ABS/RC values seem to be a consequence of the high DI_0/RC (Demetriou et al. 2007).

The photosynthetic activity of all *B. braunii* biofilms improved greatly when the cultures reached logarithmic phase. The fluorescence induction curve had a more proportional shape and lower V_J compared to those from the previous phases. The high value of V_J illustrated the actual proportion of closed reaction centres at 2 ms, resulting in less connectivity in the PS-II unit (Force et al. 2003). Other parameters, such as PI_{ABS} and F_V/F_0 , were also improved in the logarithmic phase, while ABS/RC and DI_0/RC decreased. The increases in PI_{ABS} and F_V/F_0 indicated that the algal vitality and ability to utilise light for photosynthetic activity was improving (Maxwell and Johnson 2000; Cosgrove and Borowitzka 2010). In the logarithmic phase, *B. braunii* had acclimated to the new environment, and the cells were dividing at a constant rate (Becker 1994; Basanti and Gualtieri 2014). The stationary phase begins when the algae increase their biomass during growth and the loss during the degradation process reaches an equilibrium, and one or more nutrients are depleted (Becker 1994; Basanti and Gualtieri 2014). Our study showed that a significant decrease of F_V/F_0 can be an indicator variable showing that the algae had entered stationary phase. At this stage, although the ability of *B. braunii* to utilise light for photosynthesis has reduced, the photosynthetic activity at the reaction centres was still effective (characterised by a decrease of ABS/RC and DI_0/RC). This is a consequence of the high value of PI_{ABS} , which is directly affected by the absorption of light energy and the trapping of the excited energy (Strauss et al. 2006). Further investigations should be conducted to determine the photosynthetic apparatus that is directly

affected when the algae enter the stationary phase. A decrease in V_J also occurred in the stationary phase, which indicated that the PS-II units in the photosynthesis system were still well connected (Force et al. 2003).

In conclusion, our study found that *B. braunii* has the potential to be grown as a biofilm. We also found that photosynthetic activity measurements can be used as a rapid and non-destructive way to analyse overall stress in *B. braunii* cultures during the different phases of growth. Furthermore, a visible sign in the photosynthetic parameters that shows that the algae have entered stationary phase may be an advantage in indicating the appropriate time for the lipid extraction process. However, this method still requires some improvements. Among the strains tested here, *B. braunii* BOT22 showed the highest biomass yield and lipid productivity on biofilm, which makes this strain a very good candidate for potential advanced studies. It should be noted that our long-term aim is to milk these biofilms for their hydrocarbons. The fact that *B. braunii* hydrocarbons remain in the top layers of the biofilm makes *B. braunii* biofilm cultures a very good candidate for the milking process. However, further larger scale growth studies, particularly under outdoor conditions, are necessary to indicate the reliability of *B. braunii* cultures as a promising raw material for biofuel production.

References

- Ashokkumar V, Rengasamy R (2012) Mass culture of *Botryococcus braunii* Kutz. Under open raceway pond for biofuel production. *Bioresour Technol* 104:394–399
- Banerjee A, Sharma R, Chisti Y, Banerjee U (2002) *Botryococcus braunii*: a renewable source of hydrocarbons and other chemicals. *Crit Rev Biotechnol* 22:245–279

- Basanti L, Gualtieri P (2014) Algae: anatomy, biochemistry, and biotechnology, 2nd edn. CRC Press, Boca Raton
- Becker EW (1994) Microalgae: biotechnology and microbiology. Cambridge University Press, Cambridge
- Berner F, Heimann K, Sheehan M (2015) Microalgal biofilms for biomass production. J Appl Phycol 27:1793–1804
- Borowitzka MA (2013) Energy from microalgae: a short history. In: Borowitzka MA, Moheimani NR (eds) Algae for biofuel and energy. Springer, Dordrecht, pp. 1–15
- Borowitzka MA, Moheimani NR (2013) Sustainable biofuels from algae. Mitig Adapt Strateg Glob Chang 18:13–25
- Brennan L, Blanco Fernández A, Mostaert AS, Owende P (2012) Enhancement of BODIPY505/515 lipid fluorescence method for applications in biofuel-directed microalgae production. J Microbiol Meth 90:137–143
- Brown AC, Knights BA, Conway E (1969) Hydrocarbon content and its relationship to physiological state in the green alga *Botryococcus braunii*. Phytochemistry 8:543–547
- Chaudry S, Bahri PA, Moheimani NR (2015) Pathways of processing of wet microalgae for liquid fuel production: a critical review. Renew Sust Energ Rev 52:1240–1250
- Chaudry S, Bahri PA, Moheimani NR (2017) Superstructure optimization and energetic feasibility analysis of process of repetitive extraction of hydrocarbons from *Botryococcus braunii*- a species of microalgae. Comput Chem Eng 97:36–46
- Cheng P, Ji B, Gao L, Zhang W (2013) The growth, lipid and hydrocarbon production of *Botryococcus braunii* with attached cultivation. Bioresource Technol 138:95–100
- Cirulis JT, Strasser BC, Scott JA, Ross GM (2012) Optimization of staining conditions for microalgae with three lipophilic dyes to reduce precipitation and fluorescence variability. Cytometry 81A:618–626
- Consalvey M, Perkins R, Peterson D (2005) PAM fluorescence: a beginners guide for benthic diatomists. Diatom Res 20:1–22
- Cosgrove J, Borowitzka MA (2010) Chlorophyll fluorescence terminology: an introduction. In: Suggett D, Prášil O, Borowitzka M (eds) Chlorophyll *a* fluorescence in aquatic sciences: methods and applications. Springer, Dordrecht, pp. 1–17
- Dao LH, Beardall J (2016) Effect of lead on two green microalgae *Chlorella* and *Scenedesmus*: photosystem II activity and heterogeneity. Algal Res 16:150–159
- Dayananda C, Sarada R, Usha Rani M, Shamala T, Ravishankar G (2007) Autotrophic cultivation of *Botryococcus braunii* for the production of hydrocarbons and exopolysaccharides in various media. Biomass Bioenerg 31:87–93
- de Boer K, Moheimani NR, Borowitzka MA, Bahri PA (2012) Extraction and conversion pathways for microalgae to biodiesel: a review focused on energy consumption. J Appl Phycol 24:1681–1698
- Demetriou G, Neonaki C, Navakoudis E, Kotzabasis K (2007) Salt stress impact on the molecular structure and function of the photosynthetic apparatus-the protective role of polyamines. Biochim Biophys Acta 1767:272–280
- Dixon RK (2013) Algae based biofuels. Mitig Adapt Strateg Glob Chang 18:1–4
- Eroglu E, Melis A (2010) Extracellular terpenoid hydrocarbon extraction and quantitation from the green microalgae *Botryococcus braunii* Var. Showa. Bioresour Technol 101:2359–2366
- Fon Sing S, Isdepsky A, Borowitzka MA, Moheimani NR (2013) Production of biofuels from microalgae. Mitig Adapt Strateg Glob Chang 18:47–72
- Force L, Critchley C, van Rensen J (2003) New fluorescence parameters for monitoring photosynthesis in plants: 1. The effect of illumination on the fluorescence parameters of the JIP-test. Photosynth Res 78: 17–33
- Gates M, Rogerson A, Berger J (1982) Dry to weight biomass conversion constant for *Tetrahymena elliotti* (Ciliophora, protozoa). Oecologia 55:145–148
- Govender T, Ramanna L, Rawat I, Bux F (2012) BODIPY staining, an alternative to the Nile red fluorescence method for the evaluation of intracellular lipids in microalgae. Bioresour Technol 114:507–511
- Graham LE, Graham JM, Wilcox LW (2009) Algae, 2 edn. Benjamin Cummings, San Francisco
- Heydom A, Nielsen AT, Hentzer M, Sternberg C, Givskov M, Ersbøll BK, Molin S (2000) Quantification of biofilm structures by the novel computer program COMSTAT. Microbiology 146:2395–2407
- Hillen LW, Pollard G, Wake LV, White N (1982) Hydrocracking of the oils of *Botryococcus braunii* to transport fuels. Biotechnol Bioeng 24:193–205
- Kawachi M, Tanoi T, Demura M, Kaya K, Watanabe MM (2012) Relationship between hydrocarbons and molecular phylogeny of *Botryococcus braunii*. Algal Res 1:114–119
- Lawrence JR, Neu TR, Swerhone GDW (1998) Application of multiple parameter imaging for the quantification of algal, bacterial and exopolymer components of microbial biofilms. J Microbiol Meth 32:253–261
- Liu T, Wang J, Hu Q, Cheng P, Ji B, Liu J, Chen Y, Zhang W, Chen X, Chen L, Gao L, Ji C, Wang H (2013) Attached cultivation technology of microalgae for efficient biomass feedstock production. Bioresour Technol 127:216–222
- Lu C, Vonshak A (1999) Photoinhibition in indoor *Spirulina platensis* cultures assessed by polyphasic chlorophyll fluorescence transients. J Appl Phycol 11:355–359
- Maxwell K, Johnson G (2000) Chlorophyll fluorescence—a practical guide. J Exp Bot 51:659–668
- Metzger P, Casadevall E (1989) Aldehydes, very long chain alkenylphenols, epoxides and other lipids from an alkadiene-producing strain of *Botryococcus braunii*. Phytochemistry 28: 2097–2104
- Metzger P, Largeau C (2005) *Botryococcus braunii*: a rich source for hydrocarbons and related ether lipids. Appl Microbiol Biotechnol 66:486–496
- Metzger P, Berkaloﬀ C, Casadevall E, Coute A (1985) Alkadiene- and botryococcene-producing races of wild strains of *Botryococcus braunii*. Phytochemistry 24:2305–2312
- Moheimani NR, Cord-Ruwisch R, Raes E, Borowitzka MA (2013) Non-destructive oil extraction from *Botryococcus braunii* (Chlorophyta). J Appl Phycol 25:1653–1661
- Moheimani NR, Matsuura H, Watanabe MM, Borowitzka MA (2014) Non-destructive hydrocarbon extraction from *Botryococcus braunii* BOT-22 (race B). J Appl Phycol 26:1453–1463
- Neu TR, Woelfl S, Lawrence JR (2004) Three-dimensional differentiation of photo-autotrophic biofilm constituents by multi-channel laser scanning microscopy (single-photon and two-photon excitation). J Microbiol Meth 56:161–172
- Ozkan A, Kinney K, Katz L, Berberoglu H (2012) Reduction of water and energy requirement of algae cultivation using an algae biofilm photobioreactor. Bioresource Technol 114:542–548
- Ricciardi A, Bourget E (1998) Weight-to-weight conversion factors for marine benthic macroinvertebrates. Mar Ecol Prog Ser 163: 245–251
- Roháček K (2002) Chlorophyll fluorescence parameters: the definitions, photosynthetic meaning, and mutual relationships. Photosynthetica 40:13–29
- Satpati G, Pal R (2015) Rapid detection of neutral lipid in green microalgae by flow cytometry in combination with Nile red staining-an improved technique. Ann Microbiol 65:937–949
- Schnurr PJ, Espie GS, Allen DG (2013) Algae biofilm growth and the potential to stimulate lipid accumulation through nutrient starvation. Bioresource Technol 136:337–344

- Scott SA, Davey MP, Dennis JS, Horst I, Howe CJ, Lea-Smith DJ, Smith AG (2010) Biodiesel from algae: challenges and prospects. *Curr Opin Biotechnol* 21:277–286
- Shi J, Podola B, Melkonian M (2007) Removal of nitrogen and phosphorus from wastewater using microalgae immobilized on twin layers: an experimental study. *J Appl Phycol* 19:417–423
- Strasser B, Dau H, Heinze I, Senger H (1999) Comparison of the light induced and cell cycle dependent changes in the photosynthetic apparatus: a fluorescence induction study on the green algae *Scenedesmus obliquus*. *Photosynth Res* 60:217–227
- Strasser BC, Srivastava A, Tsimilli-Michael M (2000) The fluorescence transient as a tool to characterize and screen photosynthetic samples. In: Yunus M, Pathre U, Mohanty P (eds) *Probing photosynthesis: mechanism, regulation & adaptation*. Taylor & Francis, London, pp. 445–482
- Strauss AJ, Krüger GHJ, Strasser RJ, Van Heerden PDR (2006) Ranking of dark chilling tolerance in soybean genotypes probed by the chlorophyll a fluorescence transient O-J-I-P. *Env Exp Bot* 56:147–157
- Watanabe M, Kawachi M, Hiroki M, Kasai F (2000) *Microalgae and protozoa* (trans: agency E). NIES collection list of strains, 6 edn. National Institute for Environmental Studies, Japan
- Zhang J (2013) *Culture of Botryococcus braunii*. MSc Thesis, Murdoch University, Western Australia